

Please type a plus sign (+) inside this box →

+

UTILITY
PATENT APPLICATION
TRANSMITTAL

Attorney Docket No.

PC10202A

First Named Inventor or Application Identifier

Campos, et al.

Title

FUSION PROTEINS COMPRISING CARRIERS THAT CAN
INDUCE A DUAL IMMUNE RESPONSE

Express Mail Label No.

EL162820034US

Only for new nonprovisional applications under 37 C.F.R. §1.53(b)

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

1. ☒ *Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original, and a duplicate for fee processing)

2. ☒ Specification [Total Pages 47]
(preferred arrangement set forth below)

- Descriptive title of the invention
- Cross References to Related Applications
- Statement Regarding Fed sponsored R&D
- Reference in Microfiche Appendix
- Background of the invention
- Brief Summary of the invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

3. ☒ Drawing(s) (35 U.S.C. 11.3) [Total sheets 32]

4. ☐ Oath or Declaration [Total pages]

- a. ☐ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 CFR §1.63(d))
(for continuation/divisional with Box 17 completed)
[Note Box 5 below]

- i. ☐ DELETION OF INVENTOR(S)

Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§1.63(d)(2) and 1.33(b).

5. ☐ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered to be part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)

7. Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)

- a. ☒ Computer Readable Copy
- b. ☒ Paper Copy (identical to computer copy)
- c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))

9. ☐ 37 C.F.R. §3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)

10. ☐ English Translation Document (if applicable)

11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations

12. ☒ Preliminary Amendment

13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)

14. ☐ *Small Entity ☐ Statement filed in prior application,
Statement(s) Status still proper and desired
(PTO/SB/09-12)

15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)

14. ☐ Other: Priority claimed from provisional application number 60/120,454, filed February 17, 1999.

*NOTE FOR ITEMS 1 & 14: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

17. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No: /

Prior application information:

Examiner

Group/Art Unit:

18. CORRESPONDENCE ADDRESS

☐ Customer Number or Bar Code Label (Insert Customer No. or Attach bar code label here) or ☒ Correspondence address below

Name	Paul H. Ginsburg				
Address	Pfizer Inc				
Address	235 East 42nd Street, 20th Floor				
City	New York	State	New York	Zip Code	10017-5755
Country	United States Of America	Telephone	(212)573-2369	Fax	(212)573-1939

NAME (Print/type) Kristina L. Konstas

Registration No. (Attorney/Agent)

37,864

Signature

Kristina L. Konstas

Date Feb 16, 2000

EXPRESS MAIL NO. EL162820034US

UTILITY TRANSMITTAL PTO SB 05, 3/99, (1/1)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN RE APPLICATION OF: Manuel Campos, et al. :
APPLICATION NO.: Not Yet Known : Examiner: Not Yet Assigned
FILING DATE: Herewith : Group Art Unit: Not Yet Assigned
TITLE: FUSION PROTEINS COMPRISING :
CARRIERS THAT CAN INDUCE A DUAL
IMMUNE RESPONSE
Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

PRELIMINARY AMENDMENT

Please amend the above-identified application, which is today being filed herewith, as follows:

In the Specification:

On page 1, between line 2 and line 4, please insert the following paragraph:

-- This application claims the benefit under 35 U.S.C. 119 of U.S. Provisional Patent Application No. 60/120,454, filed February 17, 1999.--

In the Claims:

In the second line of claim 11, please delete "or a transformed cell according to claim 10", and please insert --or-- before the phrase "a vector".

In the second line of claim 12, please insert --or-- before the phrase "a vector".

In the third line of claim 12, please delete "or a transformed cell according to claim 10".

In the second line of claim 13, please insert --or-- before the phrase "a vector".

In claim 13, the second and third lines thereof, please delete "or a transformed cell according to claim 10".

No fee is believed necessary for filing this Preliminary Amendment. However, should a fee be determined necessary in connection with the filing of this Preliminary Amendment, authorization is hereby given to charge such fee to Deposit Account No. 16-1455.

Respectfully submitted,

Date:

February 16, 2000

Christina L. Konkar

Kristina L. Konstas
Attorney for Applicant(s)
Reg. No. 37,864

Pfizer Inc
Patent Department, 20th Floor
235 East 42nd Street
New York, NY 10017-5755
(212) 733-6380

[illegible]

**FUSION PROTEINS COMPRISING CARRIERS THAT CAN
INDUCE A DUAL IMMUNE RESPONSE**

Field of the Invention

5 The present invention is in the field of animal and human health, and is directed to fusion proteins useful in vaccine compositions.

Background of the Invention

10 The vertebrate immune system comprises an intricate system of cells, secreted factors, and responses for protecting an organism from pathogenic infection by microbes, viruses, toxins, and other pathogens and irritants. Certain molecules, however, comprise epitopes which do not induce an effective immune response in a vertebrate because of their small size and/or because they are endogenously synthesized within the vertebrate and are therefore not perceived as "foreign" by the vertebrate's immune system. Methods for
15 producing antibodies against certain peptides which are normally non-immunogenic, such as hormones, are desirable because immunoregulation of the activity of such peptides within the organism can thereby be achieved.

20 Hormone peptides have been combined with various carrier peptides in fusion proteins to elicit an effective immune response against the hormone when an organism is vaccinated with the fusion protein. The carrier portion causes the organism's immune system to recognize and generate antibodies against the hormone peptide which it would not otherwise generate.

25 U.S. Patent 5,403,586 to Russell-Jones *et al.*, for example, relates to fusion proteins which comprise an analog of gonadotropin releasing hormone (GnRH), also known as luteinizing hormone releasing hormone (LHRH), and a TraTp analog, wherein the presence of the TraTp analog in the fusion protein helps trigger the production of anti-GnRH antibodies. TraTp is an outer membrane lipoprotein produced by certain strains of *E.coli*, as described in U.S. Patent 5,403,586, above.

30 U.S. Patent 5,422,110 to Potter *et al.* relates to carrier systems that include chimeric proteins which comprise a leukotoxin polypeptide fused to a selected antigen. The leukotoxin functions to increase the immunogenicity of the antigen. Selected antigens that are disclosed therein include GnRH, somatostatin (SRIF), and rotavirus viral protein 4 (VP4).

35 WO 90/02187 relates to fusion proteins which comprise an antigenic, hydrophilic portion, such as Hepatitis B surface antigen (HBsAg), and a peptide, such as GnRH, which alone is not substantially antigenic.

EXPRESS MAIL NO. EL162820034US

GnRH is a decapeptide endogenously produced, mainly in the hypothalamus. It is highly conserved among vertebrate species. In mammals, the GnRH gene encodes the decapeptide glu-his-trp-ser-tyr-gly-leu-arg-pro-gly with subsequent post-translational modification of the N and C termini to pyroglutamic acid and glycnamide, respectively, producing (pyro)-glu-his-trp-ser-tyr-gly-leu-arg-pro-gly-NH₂. GnRH has been shown to play a critical role in the regulation of reproductive functions in all major vertebrates by regulating the production and release of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the pituitary gland. Because FSH and LH play a role in spermatogenesis and ovulation, as well as steroidogenesis, vaccines that result in the production of antibodies against GnRH lead to the suppression of reproductive function (fertility) in both males and females, and should also control secondary sexual characteristics such as gender-related behavior. In males, LH regulates steroidogenesis in Leydig cells. Thus, active immunization of males against GnRH leads to testicular atrophy and a decrease in testosterone production and testicular function, (Ladd, A. *et al.*, 1994, Biol. Reprod. 51:1076-1083; Ladd A., 1993, Am. J. Reprod. Immunol. 29:189-194). A GnRH vaccine has been approved by the United States Food and Drug Administration as an investigational new drug for the treatment of prostate cancer (Ladd A., 1993, above). The development of a GnRH immuno-contraceptive is a useful alternative to surgical sterilization in animals, and has the added advantage of being reversible, since spermatogenesis and fertility can return to normal by simply allowing anti-GnRH titers to decline (Ladd, A. *et al.*, 1989, J. Reprod. Immunol. 15:85-101). However, since GnRH is a small self peptide and has a short half-life (WO 90/02187, March 8, 1990), it is only weakly immunogenic, even when injected with a powerful adjuvant. For example, a significant proportion of animals are not able to mount an effective antibody response against GnRH when administered in Freund's complete adjuvant. In order to generate a significant antibody response, GnRH must therefore be conjugated, chemically or recombinantly, to a carrier protein.

None of the aforementioned references, however, teach or suggest using a carrier which triggers an immunoinhibiting response against itself.

Summary of the Invention

The subject invention provides a fusion protein for producing a dual immune response in a vertebrate, which fusion protein comprises: (a) a first proteinaceous portion analogous to all or part of a peptide endogenously synthesized within the vertebrate, the activity of which peptide is to be inhibited within the vertebrate, and which proteinaceous portion by itself is incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to (b) a second proteinaceous portion analogous to all or part of an immunogen from a

pathogen, which pathogen is capable of pathogenically infecting the vertebrate; the portion (b) causing the vertebrate's immune system to recognize the portion (a) and produce a response that: (i) inhibits the activity of the peptide endogenously synthesized within the vertebrate; and (ii) protects the vertebrate from infection by the pathogen, when the vertebrate is vaccinated with an effective amount of the fusion protein.

The subject invention further provides, in a second aspect, a fusion protein for producing an immune response in a vertebrate, which fusion protein comprises: (a) a first proteinaceous portion analogous to all or part of a peptide, the activity of which peptide is to be inhibited within the vertebrate, and which proteinaceous portion by itself is incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to (b) a second proteinaceous portion analogous to all or part of a Bovine Herpes Virus Type-1 (BHV-1) antigen; the second proteinaceous portion (b) causing the vertebrate's immune system to recognize the first proteinaceous portion (a) and produce an immune response capable of inhibiting the activity of the peptide within the vertebrate when the vertebrate is vaccinated with an effective amount of the fusion protein.

The subject invention further provides fusion proteins as recited in the preceding two paragraphs which are recombinant fusion proteins.

The subject invention further provides a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the present invention.

The subject invention further provides a vector which comprises a polynucleotide molecule comprising a nucleotide sequence which encodes a fusion protein of the present invention.

The subject invention further provides a transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the present invention.

The subject invention further provides a dual-function vaccine which comprises an amount of a fusion protein as set forth above comprising: (a) a first proteinaceous portion analogous to all or part of a peptide endogenously synthesized within a vertebrate, the activity of which peptide is to be inhibited within the vertebrate, and which proteinaceous portion by itself is incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to (b) a second proteinaceous portion analogous to all or part of an immunogen from a pathogen, which pathogen is capable of pathogenically infecting the vertebrate; the portion (b) capable of causing the vertebrate's immune system to recognize the portion (a) and produce a response that: (i) inhibits the activity of the peptide endogenously synthesized within the vertebrate; and (ii) protects the vertebrate from infection by the pathogen, said fusion protein being present in the dual-function vaccine in an amount effective to inhibit the

activity of the peptide from which portion (a) is derived and to protect the vertebrate from infection by the pathogen from which portion (b) is derived, said dual-function vaccine further comprising a carrier acceptable for pharmaceutical or veterinary use.

5 The subject invention further provides a method for inhibiting the activity of an endogenously-synthesized peptide in a vertebrate and for protecting the vertebrate from a pathogenic infection, which method comprises immunizing the vertebrate with a vaccine as recited in the preceding paragraph in an amount effective to inhibit the activity of the peptide and to protect against infection by the pathogen.

10 The subject invention further provides a vaccine for inhibiting the activity of a peptide in a vertebrate which comprises a fusion protein as set forth above which comprises: (a) a first proteinaceous portion analogous to all or part of a peptide, the activity of which peptide is to be inhibited within the vertebrate, and which proteinaceous portion by itself is incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to (b) a second proteinaceous portion analogous to all or part of a BHV-1 antigen; the second proteinaceous
15 portion (b) being capable of causing the vertebrate's immune system to recognize the first proteinaceous portion (a) and to produce a response that inhibits the activity of the peptide within the vertebrate, the fusion protein being present in the vaccine in an amount effective to inhibit the activity of the peptide in the vertebrate, and the vaccine further comprising a carrier acceptable for pharmaceutical or veterinary use.

20 The subject invention further provides a method for inhibiting the activity of a peptide in a vertebrate which comprises immunizing the vertebrate with a vaccine as recited in the preceding paragraph in an amount effective to inhibit the peptide.

The subject invention further provides a method of making polyclonal antibodies directed against a peptide that is endogenously synthesized in a vertebrate which comprises
25 vaccinating such a vertebrate with an antibody-inducing amount of a fusion protein of the present invention, or a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which fusion protein comprises a portion (a) analogous to all or part of a peptide endogenously synthesized within the vertebrate; obtaining serum containing polyclonal antibodies from the vaccinated
30 vertebrate; and isolating from the serum polyclonal antibodies which bind to the endogenously-synthesized peptide; thereby making polyclonal antibodies directed against the peptide.

The subject invention further provides polyclonal antibodies directed against an endogenously-synthesized peptide made according to the method recited in the preceding
35 paragraph.

The subject invention further provides a method of making a monoclonal antibody directed against a peptide that is endogenously synthesized in a vertebrate which comprises vaccinating such a vertebrate with an antibody-inducing amount of a fusion protein of the present invention, or vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which fusion protein comprises a portion (a) analogous to all or part of a peptide endogenously synthesized within the vertebrate; and isolating a spleen cell from the vaccinated vertebrate which spleen cell excretes a monoclonal antibody that specifically binds to the endogenously-synthesized peptide; thereby making a monoclonal antibody directed against the peptide.

The subject invention further provides monoclonal antibodies directed against an endogenously-synthesized peptide made according to the method recited in the preceding paragraph.

Brief Description of the Figures

FIGURE 1: Constructs of gD/GnRH fusions: fusion proteins constructed according to the subject invention are depicted. gD in these constructs is mature (the signal sequence has been removed) and truncated (the transmembrane domain and remaining 3' sequence has been removed). GnRH is in tetrameric form.

FIGURE 2: A GnRH-tetramer clone, constructed by fusing the C termini of annealed GnRH oligonucleotides (set forth in SEQ ID NOS: 7 and 8) to the GnRH sequence (annealed oligonucleotides set forth in SEQ ID NOS: 9 and 10) in dimer clone 98BS/GnR. Flanking sequences from plasmid pBS KS+ (Stratagene) and cloning sites therein, are also depicted. This nucleotide sequence is set forth in SEQ ID NO: 14. The encoded amino acid sequence (SEQ ID NO: 15) is also shown.

FIGURE 3 (Fig.3A-3C): Nucleotide sequence (SEQ ID NO: 16) encoding BHV-1 gD within clone FlgD/Pots207nco(#79), as well as the encoded polyaminoacid gD sequence (SEQ ID NO: 17). Nucleotides 3-56 encode the signal sequence; nucleotides 1092-1169 encode the transmembrane domain. Nucleotides 57-1259 encode mature gD, and nucleotides 57-1076 encode truncated mature gD. "Gly" represents regions of glycosylation. Vector sequences flanking the gD coding sequence are shown.

FIGURE 4 (Fig.4A-4C): Alignment report (DNA alignment) comparing BHV-1 gD from clone FlgD/Pots207nco(#79) (gD/Pots, top sequence) and BHV-1 gD having GenBank Accession No. M59846 (bottom sequence) (Tikoo *et al.*, 1990, above) (GenBank DNA sequence database of the U.S. National Center for Biotechnology Information (NCBI, Bethesda, Maryland)). Clustal method with weighted residue weight table was used for this report. "TM" stands for transmembrane domain. Boxed residues in the

FlgD/Pots207nco(#79) clone are those that differ from residues in M59846. M59846 DNA is SEQ ID NO: 18.

FIGURE 5: Amino acid alignment between gD/Pots (bottom sequence) and M59846 (top sequence). Clustal method with PAM250 residue weight table was used. Residues in gD/Pots which differ from residues in M59846 are boxed. M59846 is SEQ ID NO: 19.

FIGURE 6: (Fig.6A-6C): pQE-tmgD. Nucleotide coding sequence for the tmgD, flanked by plasmid pQE-31 sequences, including a sequence encoding a 6XHIS tag, which is expressed connected to the tmgD (SEQ ID NO: 20). The amino acid sequence of the tmgD with the connected 6XHIS tag is also shown (SEQ ID NO: 21).

FIGURE 7 (Fig.7A-7C): Nucleotide coding sequence and flanking sequences for plasmid pQE-GnRH:gD (SEQ ID NO: 22). Amino acid sequence of the 4GnRH-tmgD fusion protein, including a 6XHIS tag, is also shown (SEQ ID NO: 23).

FIGURE 8 (Fig.8A-8C): pQE-gD:GnRH. Nucleotide coding sequence and plasmid flanking sequences are shown (SEQ ID NO: 24). The amino acid sequence of the tmgD-4GnRH, with a 6XHIS tag, is also shown (SEQ ID NO: 25).

FIGURE 9 (Fig.9A-9C): pQE-GnRH:gD:GnRH. Nucleotide coding sequence and plasmid flanking sequences are shown (SEQ ID NO: 26). The amino acid sequence of the 4GnRH-tmgD-4GnRH, with a 6XHIS tag, is also shown (SEQ ID NO: 27).

FIGURE 10: Comparison of expression products from bacterial vector pQE constructs. "A" is pQE-tmgD, "B" is pQE-gD:GnRH, "C" is pQE-GnRH:gD, and "D" is pQE-GnRH:gD:GnRH. The amino acids which link the gD portions, the GnRH tetramers, and the 6XHIS tags are depicted in this figure.

FIGURE 11 (Fig. 11A-11B): Nucleotide sequence (SEQ ID NO: 28) from plasmid pCMV-tgD encoding a truncated gD, and deduced amino acid sequence (SEQ ID NO: 29) of the truncated gD expression product including the signal sequence.

FIGURE 12 (Fig. 12A-12B): Nucleotide sequence (SEQ ID NO: 30) from plasmid pCMV-gD:GnRH (ATCC Accession No. 203370) encoding a tgD-4GnRH fusion protein, with deduced amino acid sequence (SEQ ID NO: 31) of the fusion protein product including signal sequence.

Detailed Description of the Invention

Fusion Proteins

In a first aspect, the subject invention provides fusion proteins that induce in a vertebrate a dual immune response that both inhibits the activity of a peptide endogenously synthesized by the vertebrate and also inhibits a pathogenic infection in the vertebrate. The inhibition of the endogenously-synthesized peptide is obtained by connecting a first

proteinaceous portion which is analogous to all or part of the endogenously-synthesized peptide to a carrier, the carrier being a proteinaceous portion analogous to all or part of an immunogen from a pathogen capable of pathogenically infecting the vertebrate. In addition to functioning as a carrier (*i.e.* enhancing an immune response against the analog of the endogenously-synthesized peptide or part thereof), the portion analogous to the immunogen or immunogen part from the pathogen also induces a response against itself in the vertebrate and thus protects the vertebrate from infection by the pathogen.

Since two major causes of economic loss in feedlot cattle in the United States and in range-fed cattle globally are bovine respiratory disease (BRD) caused by BHV-1 infection and sexual and aggressive behavior, a product that will simultaneously treat BRD and also inhibit secondary sexual characteristics, *e.g.*, aggression, would improve carcass quality and productivity by eliminating or reducing these infectious and endocrine causes of production losses in cattle. Thus, as one embodiment of the present invention, glycoprotein D (gD), which is an immunogen from BHV-1, was selected as a carrier to combine with a GnRH peptide in a fusion protein in order to regulate GnRH activity in cattle while simultaneously providing protection against BRD.

Certain proteinaceous portions that are analogs to an immunogen from a pathogen or part of an immunogen from a pathogen, such as the BHV-1 glycoprotein analogs described herein, have not previously been disclosed or suggested as carriers. Thus, a second aspect of the subject invention provides a fusion protein for producing an immune response in a vertebrate, which fusion protein comprises as a carrier a proteinaceous portion analogous to all or part of a BHV-1 antigen. In this second aspect, the vertebrate need not be a vertebrate which is capable of being pathogenically infected by BHV-1; the BHV-1 antigen analog simply acts as a carrier that induces an immune response inhibiting the activity of proteinaceous portion (a).

Thus, if a fusion protein of this invention comprises, for example, a portion (a) analogous to all or part of a GnRH peptide, and a portion (b) analogous to all or part of a BHV-1 antigen, such a fusion protein will produce a dual immune response in cattle, but will also be useful in other vertebrate species for inhibiting GnRH activity without protecting against BHV-1 infection, as such other species are not pathogenically infected by BHV-1.

For purposes of this invention, "fusion protein" means a molecule comprising a plurality of proteinaceous portions connected together. Thus, fusion proteins of this invention include chemical conjugates (chemically connected portion (a) and (b)) and recombinant fusion proteins. A fusion protein according to this invention comprises a proteinaceous portion (a) and a proteinaceous portion (b), by which is meant that the molecule may comprise at least one portion (a) and at least one portion (b), but can comprise more than one portion

(a) and/or more than one portion (b). The portions (a) and (b) can be connected linearly. If multiple portions of (a) and/or (b) are present, then the portions can be connected linearly, or they can be connected in a branched manner, for example with one of the portions (a) or (b) centrally located in the molecule and with other portions (a) or (b) multiply connected to the central portion. Other patterns of connection which can be ascertained by those of ordinary skill in the art are included within the subject invention, as long as at least one portion (a) and one portion (b) are present in the fusion proteins.

The portions (a) and (b) can be positioned with respect to one another so as to optimize an effective immune response against the portion (a), as well as the portion (b) if desired. Such positioning can be ascertained by methods known to those of ordinary skill in the art. For example a fusion protein as described herein can be tested by Western blot with antibodies against (a) and/or (b) to determine if portions (a) and/or (b) are positioned so as to optimize binding of antibodies specific thereto.

The portions of the subject fusion proteins can be connected by means including chemical connections and recombinant connections. A "chemical connection" involves creating a chemical intermediate from one proteinaceous portion, and reacting the intermediate with another proteinaceous portion. For example, a "chemical connection" can involve forming a direct covalent bond between an organic group of one proteinaceous portion, such as portion (a), and an organic group of the other proteinaceous portion, e.g. portion (b), provided the portions have organic groups which are able to react under appropriate reaction conditions to form such a covalent bond. As another example, one of the proteinaceous portions, such as portion (a), can be derivatized to form an intermediate that contains substituents that will react with (b) portions. A "recombinant connection" involves ligating a nucleic acid encoding one proteinaceous portion to a nucleic acid encoding another proteinaceous portion, and expressing a protein therefrom in an appropriate expression system. Chemical connections and recombinant connection are known in the art and are described in further detail herein.

The term "carrier" as used herein (except when in the phrase "pharmaceutically acceptable carrier", "carrier acceptable for pharmaceutical or veterinary use", and like phrases, or as otherwise indicated) means a molecule which elicits or enhances an immune response against a second molecule when connected thereto.

The term "analogous to" as used herein to describe portions of a fusion protein, unless otherwise indicated, means "having the same or substantially the same structure as", for example, having the same or substantially the same amino acid sequence. For example, a proteinaceous portion which is analogous to a peptide endogenously synthesized by a vertebrate has the same or substantially the same amino acid sequence as the

endogenously-synthesized peptide. "Substantially the same amino acid sequence" means a polyaminoacid sequence otherwise having the amino acid sequence of the endogenously synthesized peptide, but in which one or more amino acid residues have been deleted, added, or substituted with a different amino acid residue, where the resulting polyaminoacid molecule is useful in practicing the present invention. A polyaminoacid molecule is useful in practicing the present invention if it can result in a specific immune response when in the fusion protein product. Amino acid substitution will preferably be conservative substitutions which are well-known in the art. Rules for making such substitutions include those described by Dayhof, M.D., 1978, Nat. Biomed. Res. Found., Washington, D.C., Vol. 5, Sup. 3, among others.

When a portion (a) or portion (b) of a fusion protein of the present invention is referred to herein as being "derived from" a peptide or pathogen, this means that the portion is analogous to all or part of the peptide or all or part of an immunogen (or antigen) from the pathogen, respectively.

"Part of" a peptide, antigen, or immunogen for purposes of this invention, unless otherwise indicated, is any part such that the resulting polyaminoacid molecule is useful in practicing the present invention. This means that the part must be sufficient to elicit an immune response against the pathogen from which (b) is derived and/or the peptide from which (a) is derived. Ascertaining such parts is within the ordinary skill in the art. In a preferred embodiment, the part of the peptide, antigen or immunogen comprises at least 60%, more preferably 70%, and even more preferably at least 90% of the amino acid sequence of the particular peptide, antigen or immunogen. The actual percentage of the peptide, antigen, or immunogen is less important than is including in the part those amino acid residues which will elicit an immune response against (b) and/or (a).

The terms "immunogen" and "antigen" as used herein mean a molecule which is able to trigger an effective immune response in a particular vertebrate or vertebrate species. Immunogens useful for the subject invention are proteinaceous molecules, i.e., molecules comprised of a sequence of amino acids, but which can also include non-protein groups, e.g., carbohydrate moieties.

The term "immune response" for purposes of this invention means the production of antibodies and/or cells (such as T lymphocytes) that are directed specifically or indirectly against, or assist in the decomposition or inhibition of, a particular epitope or particular epitopes. An "effective immune response" is an immune response that, regarding portion (a), is directed against one or more epitopes so as to inhibit the activity of a peptide endogenously synthesized in the vaccinated vertebrate; and, regarding portion (b), is directed against one or more epitopes of a pathogen so as to protect against the pathogen in the vaccinated vertebrate. "Triggering an immune response" and like phrases as used herein mean inducing

and/or enhancing an immune response in a vertebrate in response to vaccination. Phrases such as "inhibition of infection" and "protection from infection" refer not only to the absolute prevention of infection, but also to any detectable reduction in the degree or rate of infection by such a pathogen, or any detectable reduction in the severity of the disease or any symptom or condition resulting from infection by the pathogen in the vaccinated animal as compared to an unvaccinated animal. A response which inhibits infection may be induced in animals which have not previously been infected with the pathogen and/or are not infected with the pathogen at the time of vaccination. Such phrases are intended also to include inhibiting the rate or degree of infection in an animal already infected with the pathogen at the time of vaccination.

The term "dual immune response" as used herein means an effective immune response as defined above which inhibits the activity of more than one peptide, and preferably two different peptides, for example an endogenously-synthesized hormone peptide and a viral peptide.

A "dual-function vaccine", as used herein, means a vaccine which can produce an immune response in a vertebrate vaccinated therewith that is directed against more than one peptide, and preferably two different peptides, within the vertebrate, for example a hormone endogenously synthesized by the vertebrate and a viral peptide of a virus which pathogenically infects the vertebrate.

The phrase "endogenously-synthesized peptide", as used herein and unless otherwise indicated, means a peptide which is synthesized by a vertebrate as part of the vertebrate's metabolic functioning. Examples of endogenously-synthesized peptides include, but are not limited to, hormones and enzymes.

"Inhibiting the activity of a peptide" and like phrases used herein mean interfering with the peptide's ability to perform its normal function, for example its ability to catalyze a biochemical reaction (if the peptide is an enzyme), to trigger a biophysical response (if the peptide is a hormone), or to participate in viral infectivity or replication (if the peptide is a viral peptide). The phrases "amount effective to inhibit the activity of the peptide from which portion (a) is derived", "amount effective to inhibit GnRH activity", and the like, refer to that amount of fusion protein capable of inducing an immune response which is sufficient to interfere with the peptide's ability to perform its function, such as preventing GnRH from stimulating or reducing the ability of GnRH to stimulate the release of LH or FSH, or interfering with a surface protein of a virus so that it is unable to infect cells, thereby inhibiting replication and infection by the virus. An effective amount may be administered as either a single dose of a vaccine or multiple doses of a vaccine.

As used herein, the phrases "amount effective to inhibit infection by the pathogen from which (b) is derived", "amount effective to inhibit BHV-1 infection", "amount effective to protect against infection", and the like, refer to that amount of fusion protein or vaccine capable of protecting a vertebrate from infection as defined above. An effective amount may be administered as either a single dose of a vaccine or multiple doses of a vaccine.

A "vertebrate", as used herein, refers to any species having a backbone or spinal column, namely fish, amphibians, reptiles, birds, and mammals. Examples of vertebrates which can benefit from the vaccine of the subject invention include, but are not limited to, humans, chickens, pigs, dogs, cats cows, goats, sheep and horses, among others. Preferably, the vertebrate is a mammal.

The term "pathogenically infecting" as used herein refers to the ability of a pathogen to infect a vertebrate in a manner or to a degree that results in a detectable diseased condition in the vertebrate. BHV-1, for example, pathogenically infects cattle but not humans.

Peptides that can be used as a source for preparing a portion (a) of a fusion protein of the present invention include, but are not limited to the following: 1) cholecystokinen (Eng, J. *et al.*, 1990, Regul. Pept. 30(1):15-9); a fusion protein of the present invention comprising a portion (a) analogous to all or part of cholecystokinen can be used to promote appetite in a vertebrate; 2) vasoactive intestinal peptide (Nilsson, A., 1975, FEBS Lett. 60(2):322-6), inhibition of which causes a decrease in prolactin secretion which in turn discourages brooding behavior in chickens, thus resulting in increased egg production; 3) growth hormone and growth hormone fragments (Seeburg, P.H. *et al.*, 1983, DNA 2(1):37-45); a fusion protein enhancing the activity of growth hormone can promote growth in an animal; 4) growth hormone releasing hormone and fragments thereof (Gubler, U. *et al.*, 1983, Proc. Natl. Acad. Sci. U.S.A. 80(14), 4311-4314); antibodies may also enhance the growth promoting activity of growth hormone releasing hormone; 5) gastrin (Dimaline, R. *et al.*, 1986, FEBS Lett. 205(2):318-22; Kim, S.J. *et al.*, 1991, DNA Seq. 1(3):181-7; Kariya, Y. *et al.*, 1986, Gene 40(1-3):345-52) and gastrin releasing peptide (Spindel, E.R. *et al.*, 1986, Proc. Natl. Acad. Sci. USA 83(1):19-23); a fusion protein inhibiting gastrin and/or gastrin releasing peptide activity is useful, *inter alia*, in inhibiting gastric secretions, and therefore in treating ulcers; treating stomach, small intestine and/or colon cancer; and in promoting appetite; 6) IgE peptides (Batista, F.D. *et al.*, 1995, Nucleic Acids Res. 23(23):4805-11); fusion proteins inhibiting IgE are useful for alleviating and/or preventing allergies, especially allergic skin reactions; 7) an angiotensin peptide, including angiotensin peptides I, II, III, and IV (U.S. Patent 5,612,360 to Boyd *et al.*; U.S. Patent 5,599,663 to Vaughan; 5,629,292 to Rodgers and DiZerega; U.S. Patent 5,635,359 to Brunner and Nussberger); a fusion protein inhibiting the activity of an angiotensin peptide is useful for treating, e.g., hypertension in a mammal; 8)

myostatin (Kambadur, R. *et al.*, 1997, Genome Res. 7(9):910-6); inhibiting myostatin activity enhances skeletal muscle growth in an animal, without harming meat quality, and therefore can be desirable for increasing meat production in an animal; 9) inhibin or fragments thereof (U.S. Patent 5,786,179 to Kousoulas *et al.*; U.S. Patent 5,665,568 to Mason and Seeburg); a
5 fusion protein that inhibits the activity of inhibin can be used to treat infertility due to irregular production of follicle stimulating hormone in a female animal; 10) somatostatin (U.S. Patent 5,422,110, above; Shen, L.P. *et al.*, 1982, Proc. Natl. Acad. Sci. USA 79(15):4575-9; Su, C.J. *et al.*, 1988, Mol. Endocrinol. 2(3):209-16); a fusion protein inhibiting somatostatin is useful, e.g., for stimulating growth; and 11) cytokine peptides such as tumor necrosis factor (U.S.
10 Patent 5,795,967 to Aggarwal *et al.*) and interleukin-1 (Masaaki, Y. *et al.*, JP 1994073095-A 1 (March 15, 1994)); inhibiting cytokine activity in an animal can alleviate immune-potentiated inflammation, for example inflammation associated with allergies. The preceding peptides, their amino acid sequences and physiological actions, are well known in the art. The aforementioned publications describing these peptides are hereby incorporated by reference
15 in their entireties.

Examples of immunogens from which proteinaceous portions useful for portion (b) can be derived include, but are not limited to, the following immunogens: 1) OmpW (U.S. Provisional Patent Application No. 60/105,285, filed October 22, 1998; encoded by plasmid pER418 present in host cells of strain Pz418 deposited with the American Type Culture
20 Collection (otherwise known as the ATCC (Manassas, Virginia, USA) under ATCC Accession No. 98928; SEQ ID NO:44 (deduced amino acid sequence of OmpW)); OmpA1 (U.S. Provisional Patent Application No. 60/105,285, encoded by plasmid pER419 present in host cells of strain Pz419 deposited with the ATCC under ATCC Accession No. 98929; SEQ ID NO:45 (deduced amino acid sequence of OmpA1)); OmpA2 (U.S. Provisional Patent
25 Application No. 60/105,285; encoded by plasmid pER420 present in host cells Pz420 deposited with the ATCC under the designation ATCC Accession No. 98930; SEQ ID NO:46 (deduced amino acid sequence of OmpA2)); OmlA serotype 1 and serotype 5 (U.S. Patent No. 5,441,736 to Gerlach *et al.*); all from *Actinobacillus pleuropneumonia*; a proteinaceous portion analogous to all or part of OmpW, OmlA5 or OmpA can be used as a carrier in a
30 fusion protein according to the present invention while simultaneously providing swine with protection against porcine pleuropneumonia (caused by *A. pleuropneumonia* infection); 2) hepatitis B surface antigen (Hsiung *et al.*, 1984, J. Mol. Appl. Gen. 2:497); a proteinaceous portion analogous to all or part of a hepatitis B surface antigen can be used as a carrier in a fusion protein of the present while at the same time providing protection in humans against
35 hepatitis B infection; 3) an RTX ("repeat in toxin") toxin from *Actinobacillus pleuropneumonia* (Frey, J. *et al.*, 1991, Infect. Immun. 59(9), 3026-32); a proteinaceous portion analogous to all

or part of an RTX toxin as a carrier in a fusion protein of the present invention can simultaneously provide immunoprotection against *Actinobacillus pleuropneumonia* in swine and cattle; 4) β subunit of *E. coli* heat labile enterotoxin (Leong, J. *et al.*, 1985, Infect. Immun. 48(1):73-7; Inoue, T. *et al.*, 1993, FEMS Microbiol. Lett 108(2):157-61); a portion analogous to
5 all or part of β subunit of *E. coli* heat labile enterotoxin can serve as a carrier that also provides immunoprotection against *E. coli* in swine and cattle; 5) *E. coli* antigens K88 pilus or K99 pilus (Bakker, D. *et al.*, 1992, J. Bacteriol 174(20):6350-8; Simons, B.L. *et al.*, 1990, FEMS Microbiol. Lett 55(102):107-12); a proteinaceous portion analogous to all or part of K88 pilus antigen or K99 pilus antigen as a carrier in a fusion proteins of this invention can provide
10 protection against enteric *E. coli* disease in swine and cattle; 6) p68 antigen of *B. bronchiseptica* (WO 9115571-A 5 (October 17, 1991)); a proteinaceous portion analogous to all or part of p68 antigen can be used as a carrier in a fusion protein of the present invention and can provide protection against bordetella infection ("kennel cough" disease) in canines; 7) glycoprotein 53 from bovine viral diarrhea (BVD) virus (Fritzemeier, J. *et al.*, 1997, Arch. Virol.
15 142(7):1335-50); a portion analogous to all or part of glycoprotein 53 can serve as a carrier in a fusion protein and also provide protection from fatal mucosal disease in cattle; 8) viral proteins 1 and 2 of parvovirus (Xie, A. and Chapman, M.S., 1996, J. Mol. Biol. 264:497); a proteinaceous portion analogous to all or part of viral protein 1 or viral protein 2 from parvovirus can serve as a carrier in a fusion protein of the present invention and
20 simultaneously protect swine, dogs and cats from parvovirus infection; 9) a coronavirus spike protein (Kokubu, T. *et al.*, 1998, Journal of the Japan Veterinary Medical Association 51:252-55; Lewis, E.L., 1996, Bristol University Thesis (Bristol University (Clifton, Bristol, UK)); Britton, P. *et al.*, 1991, Virus Res. 21(3):181-98); a portion analogous to all or part of a coronavirus spike protein can be used as a carrier in a fusion protein and also provides
25 protection against Coronavirus infection in cattle, swine, dogs, and cats; 10) a bacterial iron-regulated outer membrane protein (Gerlach, G.F. *et al.*, 1992, Infect. Immunol. 60(8):3253-61; Thompson, S.A. *et al.*, 1993, Mol. Microbiol. 9(1):85-96); a portion analogous to all or part of such a membrane protein can be used as a carrier that also provides immunoprotection against *Actinobacillus pleuropneumonia* and/or meningitis in swine, cattle and poultry; 11)
30 rabies G protein (Shinichi, S. *et al.*, JP 1989171489-A 1 (July 6, 1989)); a proteinaceous portion analogous to all or part of rabies G protein can be used as a carrier in a fusion protein and will also simultaneously provide protection in cats, dogs, and wildlife against rabies; 12) *Streptococcus uberis* plasminogen activating protein (Leigh, J.A., 1993, WO 9314209); a proteinaceous portion analogous to all or part of *Streptococcus uberis* plasminogen activating
35 protein is useful as a carrier and also will provide treatment and/or protection against mastitis in dairy cows; 13) influenza virus hemagglutinin protein (Hovanec, D.L. and Air, G.M., 1984,

EXPRESS MAIL NO. 94162820034US

Virology 139(2):384-92) and influenza virus nucleocapsid protein (Lindstrom, S.E. *et al.*, 1998, J. Virol. 72(10):8021-31); a portion analogous to all or part of either of these proteins can be used as a carrier in a fusion protein of this invention and will simultaneously provide immunoprotection against influenza in humans, swine, and poultry; 14) tetanus toxoid (Fairweather, N.F. *et al.*, 1986, J. Bacteriol. 165(1):21-7; Niemann, H., 1986, EMBO J. 5(10):2495-502); a proteinaceous portion analogous to all or part of tetanus toxoid can be used as a carrier in a fusion protein that will also provide protection in humans, horses, and cattle against tetanus; 15) pertussis toxoid (Nicosia, A. *et al.*, 1986, Proc. Natl. Acad. Sci. USA 83(13):4631-5); a proteinaceous portion analogous to all or part of pertussis toxoid can serve as a carrier in a fusion protein and will provide immunoprotection against pertussis in humans; 16) a herpes virus glycoprotein (Gompels, U.A. *et al.*, 1992, DNA Seq. 3(1):25-39; Misra, V. *et al.*, 1988, Virology 166:542-9; Whitbeck, J.C., *et al.*, 1988, J. Virol. 62:3319-27; Fitzpatrick, D.R. *et al.*, 1989, Virology 173:46-57); a proteinaceous portion analogous to all or part of a herpes virus glycoprotein can serve as a carrier in a fusion protein of this invention and can function also in the fusion protein to provide immunoprotection from herpes in humans and cattle; 17) enterohemorrhagic *E. coli* intimin protein (Jerse, A.E. *et al.*, 1990, Proc. Natl. Acad. Sci. USA 87(20):7839-43); a portion analogous to all or part of enterohemorrhagic *E. coli* intimin protein can function as a carrier and also provide protection against hemorrhagic disease in species including humans and cattle; 18) VP2 (Cao, Y.C. *et al.*, 1995, Ping Tu Hsueh Pao 11(3):234-41); a portion analogous to all or part of VP2 can function as a carrier and can also provide immunoprotection against infectious bursa disease in poultry; and 19) F and G proteins of respiratory syncytial virus (Schrijver, R.S. *et al.*, 1997, Archives of Virology 142(11):2195-2210; Furze, J.M. *et al.*, 1997, Virology 231(1):48-58); a proteinaceous portion analogous to all or part of F protein or G protein can act as a carrier and will also provide immunoprotection against Bovine Respiratory Syncytial Virus in cattle. The preceding immunogens and their amino acid sequences are known in the art. The aforementioned publications describing the preceding immunogens are hereby incorporated by reference in their entireties.

Different proteinaceous portions (a) and (b), each portion analogous to all or part of a peptide or immunogen described in one of the preceding paragraphs or another known peptide or immunogen, can be combined according to the present invention to form a fusion protein specifically designed for a particular vertebrate, e.g., a cow, pig, chicken, or human, or a particular category of vertebrates, e.g., mammals or primates, to inhibit the activity of a particular peptide in the vertebrate while simultaneously protecting the vertebrate from infection by a certain pathogen.

As an example, GnRH is a reproductive system hormone synthesized by cattle. Inhibiting GnRH activity in cattle will provide a beneficial reduction in expression of sexual characteristics such as aggressive behavior. Since BHV-1 pathogenically infects cattle, an immunogen from BHV-1 can be used as a carrier with GnRH. Thus, in one embodiment, a portion (a) analogous to all or part of a GnRH peptide and a portion (b) analogous to all or part of an immunogen from BHV-1 are connected to provide a fusion protein that induces a dual immune response in cattle that both inhibits GnRH activity and protects against BHV-1 infection.

In another non-limiting example, the subject invention provides a fusion protein wherein portion (a) is analogous to all or part of a growth hormone, and wherein portion (b) is analogous to all or part of a BHV-1 antigen. Such a fusion protein is useful to regulate growth in cattle while providing a protective immune response against BHV-1.

In another example, portion (a) is analogous to all or part of an IgE peptide and portion (b) is analogous to all or part of p68 antigen of *B. bronchiseptica*. The resulting fusion protein is useful for treating or preventing allergies, especially allergic skin reactions, in dogs while providing a protective immune response against bordetella.

In still another example, portion (a) is analogous to all or part of cholecystokinin and portion (b) is analogous to all or part of OmpW, OmlA serotype 1, OmlA serotype 5, Omp A1, or OmpA2 from *Actinobacillus pleuropneumonia*. Such a fusion protein is useful for encouraging appetite in swine while simultaneously providing a protective immune response against porcine pleuropneumonia.

The proteinaceous portions (a) and (b) for the fusion proteins of the invention can be obtained according to methods known in the art. For example, either or both of portion (a) or portion (b) can be obtained by purification from natural sources. Alternatively, either or both of portion (a) or portion (b) can be obtained by synthetically linking amino acids together. Alternatively, either or both of portion (a) or portion (b) can be recombinantly synthesized using well-known recombinant techniques from a polynucleotide molecule comprising a nucleotide sequence encoding the portion (a) or the portion (b). Preferably, a polynucleotide molecule comprising a nucleotide sequence encoding portion (a) is ligated to a polynucleotide molecule comprising a nucleotide sequence encoding portion (b), so that the entire fusion protein is synthesized recombinantly.

Recombinant techniques within the ordinary skill in the art can be utilized to prepare polynucleotide molecules that encode portions (a) and (b) of the subject fusion proteins. Such techniques are described, among other places, in Maniatis, *et al.*, 1989, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Ausubel, *et al.*, 1989, Current Protocols in Molecular Biology, Greene Publishing Associates & Wiley

Interscience, NY; Sambrook, *et al.*, 1989, Molecular Cloning: A Laboratory Manual, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Innis *et al.*, (eds), 1995, PCR Strategies, Academic Press, Inc., San Diego; and Erlich (ed), 1992, PCR Technology, Oxford University Press, New York, all of which are incorporated herein by reference.

5 The amino acid sequences of many hormone peptides are well known in the art. Some known hormone peptides are described above. As another example, the amino acid sequence of GnRH is known in the art (see, *e.g.*, Ladd, A., 1993, above). The amino acid sequence of GnRH is also provided herein (SEQ ID NO: 13). Alternatively, if the amino acid sequence of a hormone is not known, it may be determined using standard techniques, such
10 as by performing repeated Edman degradation cycles on a purified protein fraction followed by amino acid analysis using HPLC (high pressure liquid chromatography) (see, *e.g.*, U.S. Patent 5,422,110, above). Likewise, a proteinaceous portion which is the same or substantially the same as an immunogen from a pathogen can be obtained according to standard techniques, from a known amino acid sequence or by ascertaining the amino acid
15 sequence as described above. A proteinaceous portion that is substantially the same as an immunogen from a pathogen can be determined, for example, by comparing the amino acid content of the proteinaceous portion to the known amino acid content of the immunogen, or by performing a sequence alignment comparing the proteinaceous portion to the immunogen amino acid sequence, using known techniques.

20 Examples of BHV-1 antigens from which proteinaceous portion (b) can be derived include, but are not limited to, BHV-1 gB, BHV-1 gC, and BHV-1 gD (also known in the art as BHV-1gI, gIII and gIV, respectively). Methods for obtaining proteinaceous portions which are analogous to all or part of such antigens are described above. For example, U.S. Patent 5,151,267 to Babiuk *et al.* discloses the nucleotide sequences and deduced amino acid
25 sequences of BHV-1 gI, gIII, and gIV. See, also, U.S. Patent 5,585,264 to Babiuk *et al.* In addition, U.S. Patent 5,545,523 to Batt *et al.* discloses BHV-1-specific oligonucleotides useful in the amplification of BHV-1 gI and gIV gene sequences. Furthermore, methods of purifying BHV-1 glycoproteins from virus-infected cell cultures have been described (Babiuk, L.A. *et al.*, 1987 Virology 159:57-66). The amino acid sequence of full length BHV-1 gD as published in
30 Tikoo *et al.*, 1990, above, is provided herein (see Figure 5 and SEQ ID NO: 19). Expression of full length mature BHV-1 gD has been performed in baculovirus, adenovirus, vaccinia virus and *E. coli* systems (van Drunen Littel-van den Hurk, S. *et al.*, 1993, Vaccine 11:25-35). The disclosures and teachings of the aforementioned patents and publications are incorporated herein by reference. Another example of a BHV-1 gD antigen which is useful, in whole or in
35 part, for a fusion protein of the subject invention is the BHV-1 gD polyaminoacid encoded by clone, FlgD/Pots207nco(#79) (see Figure 3 and SEQ ID NO: 17).

Although any part of a BHV-1 antigen which is able to stimulate an immune response that inhibits the peptide from which portion (a) is derived and, as in the first aspect of the invention, an immune response that protects a cow from BHV-1 infection can be used in the fusion proteins of this invention, examples of preferred parts of BHV-1 gD which can be used in this invention are truncated gD (tgD), mature gD (mgD), and truncated mature gD (tmgD). Truncated gD (tgD) refers to a gD protein wherein the transmembrane domain, optionally with downstream and/or upstream nucleotides, has been completely or partially removed. The transmembrane domain for gD is known in the art as a particular polyaminoacid region of gD of generally highly hydrophobic amino acids. The transmembrane domain of gD/Pots, BHV-1 gD encoded by clone FlgD/Pots207nco(#79), is depicted in Figure 3. The transmembrane domain for gD/Pots starts at amino acid 364 (valine) and ends at amino acid 389 (tyrosine). Mature gD refers to a gD protein which has no signal sequence at the amino-terminal end. The signal sequence of full length gD/Pots is depicted in Figure 3. In another embodiment, the proteinaceous portion (b) of the fusion protein of the subject invention can comprise a heterologous signal sequence attached to the amino terminal end of the protein. Alternatively, portion (b) can comprise no signal sequence. In one embodiment of the invention, portion (b) is analogous to a BHV-1 gD which is both truncated and mature (tmgD). An example of a truncated mature gD antigen is provided in SEQ ID NO: 35. An example of truncated gD antigen that is not mature is provided in SEQ ID NO: 29.

As used herein, "tgD" refers to a BHV-1 gD protein which is truncated as described above, "mgD" refers to a BHV-1 gD protein which is mature as described above, and "tmgD" refers to a BHV-1 gD protein which is both truncated and mature.

The term "GnRH peptide" means, unless otherwise indicated, a molecule having the amino acid sequence of SEQ ID NO: 13. In one embodiment, the subject fusion proteins comprise multiple portions (a) analogous to a GnRH peptide. In preferred embodiments, the fusion proteins of this invention comprise one or more portions analogous to four GnRH peptides consecutively linked, *i.e.*, one or more portions analogous to a GnRH tetramer. In a preferred embodiment, a fusion protein of the present invention comprises a 4GnRH portion. As used herein, "4GnRH" refers to a GnRH tetramer having four GnRH peptides consecutively linked in the same amino-terminal/carboxy-terminal orientation. Preferably, the fusion proteins of the subject invention comprise one or more GnRH tetramers, each tetramer having the amino acid sequence shown in SEQ ID NO: 15.

Hyphenated expressions provided herein and containing the terms "4GnRH", "tmgD", "tgD", and "mgD" (as defined above) indicate fusion proteins which comprise polyaminoacid portions corresponding to the terms linked from left to right in the order indicated, the left end corresponding to the amino terminal end of the fusion protein and the right end corresponding

to the carboxy terminal end of the fusion protein. The polyaminoacid portions can be directly linked to one another or they can be linked indirectly, *i.e.* the portions can be separated by one or more (for example from 1 to 10, preferably from 1 to 3) amino acids. Thus, "tmgD-4GnRH" refers to a fusion protein having a truncated mature gD portion connected, directly or indirectly, to a 4GnRH portion, the carboxy terminal end of the truncated mature gD portion being linked (directly or indirectly) to the amino terminal end of the 4GnRH. As another example, "tgD-4GnRH" refers to a fusion protein having the amino terminal end of 4GnRH portion connected to the carboxy terminal end of a truncated gD antigen which is not mature. As another example, "4GnRH-tmgD-4GnRH" refers to a 4GnRH portion having a carboxyl end linked to the amino end of a tmgD portion, which tmgD portion in turn is linked by its carboxyl end to the amino end of a second 4GnRH portion. Fusion proteins of the subject invention include, but are not limited to, the examples of fusion proteins described in this paragraph. Another example of a fusion protein of this invention is tmgD-4GnRH. In any of the aforementioned examples, the portions can be linked directly or indirectly.

As discussed above, proteinaceous portions (a) and (b) can be connected chemically by means of chemical linkers and techniques which are well known in the art. As an example, certain amino acids on a portion (a) or (b), for example on a gD analog (b) portion, may be chemically activated with a reagent, such as iodoacetamide. Remaining portions (a) or (b), for example GnRH monomers and/or multimers, may be added. In this example, terminally incorporated cysteine residues on GnRH react with activated lysine residues on the gD analog. This reaction results in fusion proteins according to the subject invention which comprise a central gD analog portion having multiple GnRH analogs connected thereabout at several lysine residues. In another example, portions (b) analogous to a BHV-1 antigen may be combined together with portions (a) analogous to GnRH monomers or multimers in the presence of ethyl-dimethylaminopropylcarbodiimide (EDAC) and N-hydroxy succinimide (NHS) (see Bernatowics, M. and Matsueda, G., 1986, Analytical Biochemistry 155:95-102). This reaction also results in a central portion (b) analogous to all or part of a BHV-1 antigen with multiple portions (a) analogous to GnRH monomers or multimers chemically connected thereabout. The chemically synthesized fusion proteins of the present invention can also optionally be chemically modified to comprise substituents other than amino acids, for example carbohydrate substituents, using known techniques. Other chemical techniques for combining proteinaceous portions, either with multiple attachments to a proteinaceous center or linear linkages of proteinaceous portions, can be used to chemically synthesize fusion proteins of the present invention using known techniques. Techniques for preparing chemically-synthesized fusion proteins of the present invention are described, among other

places, in Dunn and Pennington, 1994, Methods in Molecular Biology, Vol. 26, Chap. 10 (Humana Press Inc.), which is incorporated herein by reference.

The subject invention also provides recombinant fusion proteins as described above. Examples of recombinant fusion proteins according to the present invention include the
5 recombinant fusion protein encoded by the plasmid pCMV-gD:GnRH the plasmid pQE-gD:GnRH, the recombinant fusion protein encoded by the plasmid pQE-GnRH:gD:GnRH, and the recombinant fusion protein encoded by the plasmid pQE-GnRH:gD. Cells containing these plasmids have been deposited with the American Type Culture Collection (ATCC Manassas, Virginia, USA); they have been assigned accession numbers 203370, 98953,
10 98955, and 98954, respectively. Another example of a recombinant fusion protein of the subject invention, is the recombinant fusion protein expressed by the baculovirus construct Bac-gD:GnRH. Bac-gD:GnRH has also been deposited with the ATCC and has been assigned ATCC accession number VR-2633. The aforementioned pQE plasmids and baculovirus construct are particularly useful for *in vitro* expression of fusion proteins. The
15 plasmid pCMV-gD:GnRH is particularly useful for *in vivo* expression.

Recombinant fusion proteins according to this invention may optionally comprise portions which assist in purifying the fusion proteins from the reaction medium pursuant to *in vitro* transcription and translation. An example of a polyaminoacid sequence which can assist in purification of a recombinant protein from the medium is a 6XHIS tag. The phrase "6XHIS
20 tag" is used interchangeably in this application with "6XHIS leader". The sequence of the 6XHIS tag encoded by the vector pQE-31 is provided in SEQ. ID NO: 37. Proteins comprising a 6XHIS tag can be purified from the media by passing the media through a nickel column such as Ni-NTA column from Qiagen (Chatsworth, CA). Another example of a portion that can assist in purifying recombinant fusion proteins of this invention pursuant to *in vitro*
25 expression is the FLAGTM epitope tag (International Biotechnologies Inc., New Haven, CT) which is a hydrophilic marker peptide. The gene encoding the FLAGTM epitope tag can be inserted by standard techniques into a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of this invention at a point corresponding, e.g., to the amino or carboxyl terminus of the fusion protein. A fusion protein expressed therefrom can
30 then be detected and affinity-purified using commercially available anti-FLAGTM antibodies.

Other means of purifying recombinant proteins expressed *in vitro* are well-known in the art and can be used to purify the recombinant fusion proteins of the subject invention. Such methods are described, among other places, in Marshak, D.R., *et al.*, 1996, Strategies for Protein Purification and Characterization: a Laboratory Course Manual, Cold Spring
35 Harbor Laboratory Press, Cold Spring Harbor, NY.

EXPRESS MAIL NO. 92162820034US

Once a fusion protein of the present invention has been obtained, it can be characterized if desired by standard methods, including by SDS-PAGE, size exclusion chromatography, amino acid sequence analysis, etc. The fusion protein can be further characterized using hydrophilicity analysis (see, e.g., Hopp and Woods, 1981, Proc. Natl. Acad. Sci. USA 78:3824), or analogous software algorithms, to identify hydrophobic and hydrophilic regions. Structural analysis can be carried out to identify regions of the fusion protein that assume specific secondary structures. Biophysical methods such as X-ray crystallography (Engstrom, 1974, Biochem. Exp. Biol. 11: 7-13), computer modeling (Fletterick and Zoller (eds), 1986, in: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), and nuclear magnetic resonance (NMR) can be used to map and study potential sites of interaction between the polypeptide and other putative interacting proteins/receptors/molecules such as antibodies.

Polynucleotide Molecules and Vectors Encoding Fusion Proteins

The subject invention further provides a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the present invention. Examples of such polynucleotide molecules include, but are not limited to, a polynucleotide molecule comprising the nucleotide sequence set forth in SEQ ID NO: 34, which encodes a 4GnRH-tmgD fusion protein; a polynucleotide molecule comprising the nucleotide sequence set forth in SEQ ID NO: 40, which encodes a 4GnRH-tmgD-4GnRH fusion protein; and a polynucleotide molecule comprising the nucleotide sequence set forth in SEQ ID NO:41, which encodes a tmgD-4GnRH fusion protein.

The subject invention also provides cloning and expression vectors comprising a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the invention. The term "vector", as used herein, means a unit comprising genetic information (in the form of polynucleotide sequences), which information is able to express polyaminoacids and/or program the replication of the unit when appropriate conditions and resources (e.g. amino acids, nucleotides, and transcription factors) are present. Examples of such units include viruses, plasmids, and cosmids.

As used herein, the terms "nucleotide sequence", "coding sequence", "polynucleotide", "polynucleotide sequence", and the like, refer to both DNA and RNA sequences, which can either be single-stranded or double-stranded, and can include one or more prokaryotic sequences, eukaryotic sequences, cDNA sequences, genomic DNA sequences, including exons and introns, and chemically synthesized DNA and RNA sequences.

EXPRESS MAIL NO. EL16282003405

Production and manipulation of polynucleotide molecules of the subject invention comprising nucleotide sequences encoding portions (a) and (b) of the subject fusion proteins are within the ordinary skill in the art and can be carried out according to recombinant techniques described, among other places, in Maniatis, et al., above; Ausubel, *et al.*, above; 5 Sambrook, et al., above; Innis et al., above; and Erlich, above. Nucleotide sequences encoding many hormone peptides and viral antigen peptides are known in the art, and such information can be used to prepare coding regions for the proteinaceous portions (a) and (b). Such sequences are provided, among other places, in the references cited above describing immunogens and peptides useful in the present invention. Alternatively, the nucleotide 10 sequences of peptides and viral antigens can be deduced using known methods in molecular biology.

Nucleotide sequences encoding portion (a) and/or portion (b) can be synthetically prepared. The desired sequence can be prepared from overlapping oligonucleotides. See, e.g., Edge, 1981, *Nature* 292:756; Nambair *et al.*, 1984 *Science* 223:1299; Jay *et al.*, 1984, *J.* 15 *Biol. Chem.* 259:6311; and U.S. Patent 5,422,110, above.

As another example, the amino acid sequence of a peptide or antigen can be used to design probes for identifying the gene encoding the peptide or antigen in a genomic library. In this method, oligonucleotide probes are prepared encoding a portion of the amino acid sequence of the peptide or antigen. The oligonucleotide probes are used to screen a suitable 20 DNA library for genes encoding the peptide or the antigen. Generally, the DNA library which is screened is a library prepared from genomic DNA or genomic RNA (cDNA) from an appropriate source, such as from a cell or tissue expressing the peptide or from a virus encoding the antigen. Techniques for isolating genes in this manner are well-known in the art.

Nucleotide sequences homologous to sequences obtained as described herein to 25 encode immunogens or peptides can also be utilized in the present invention. For purposes of the subject invention, a second nucleotide sequence is "homologous" to a first nucleotide sequence when it encodes the same protein, peptide, or other polyaminoacid as the first nucleotide sequence, or when it encodes a polyaminoacid that is sufficiently similar to the polyaminoacid encoded by the first nucleotide sequence so as to be useful in practicing the 30 present invention. Since the genetic code is degenerate, a homologous nucleotide sequence can include any number of "silent" base changes, *i.e.* nucleotide substitutions that nonetheless encode the same polyaminoacid. A homologous nucleotide sequence can further contain non-silent mutations, *i.e.* base substitutions, deletions, or additions resulting in amino acid differences in the encoded polyaminoacid, so long as the sequence of the 35 polyaminoacid remains useful for practicing the present invention. A second nucleotide sequence that is homologous to a first nucleotide sequence is preferably one that hybridizes

to the complement of the first nucleotide sequence under moderately stringent conditions, *i.e.*, hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.2xSSC/0.1% SDS at 42°C (see Ausubel *et al.*, above). More preferably, homologous nucleotide sequences hybridized to one another under highly stringent conditions, *i.e.*, hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% SDS, 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel *et al.*, above).

After having obtained polynucleotide molecules comprising nucleotide sequences encoding portions (a) and (b), these polynucleotide molecules can be ligated together using suitable enzymes and known techniques to form a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of this invention.

Examples of coding sequences useful in constructing polynucleotide molecules comprising sequences encoding fusion proteins of the present invention, and vectors comprising such polynucleotide molecules, include, but are not limited to, the sequence presented in SEQ ID NO: 16, which encodes the BHV-1 gD antigen FlgD/Pots expressed by clone FlgD/Pots207nco(#79), set forth in SEQ ID NO: 17; the sequence presented in SEQ ID NO: 18, which encodes M59846 BHV-1 gD, set forth in SEQ ID NO: 19; the sequence presented in SEQ ID NO: 28, which encodes a truncated gD antigen that is not mature, set forth in SEQ ID NO: 29; and the sequence presented in SEQ ID NO: 36, which encodes a truncated mature gD, set forth in SEQ ID NO: 35. An example of a nucleotide sequence that encodes a GnRH monomer is set forth in SEQ ID NO: 33. An example of a sequence which encodes a GnRH tetramer, namely the GnRH tetramer having the amino acid sequence set forth in SEQ ID NO: 15, is set forth in SEQ ID NO: 32.

In one embodiment, a vector of the subject invention is suitable for *in vitro* expression of a fusion protein, such as a plasmid which is capable of transfecting a host cell such as a bacterial cell and expressing the fusion protein in the bacterial cell. Examples of plasmid vectors include plasmids, such as recombinant pQE plasmids, capable of transfecting bacteria and expressing the fusion proteins of this invention. Examples of some prokaryotic expression vector plasmids into which a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the invention can be inserted include pQE-50 and pQE-31 (Qiagen, Chatsworth, CA), pUC8, pUC9, pBR322 and pBR 239 (Biorad Laboratories, Richmond, CA), pPL and pKK223 (Pharmacia, Piscataway, NJ). Other plasmids known in the art can also be used to prepare vectors comprising a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of this invention, and such plasmids can be ascertained by those of ordinary skill. Preferred plasmids which are capable of expressing fusion proteins of the invention *in vitro* include pQE-gD:GnRH (ATCC Accession No. 98953), pQE-GnRH:gD:GnRH (ATCC Accession No. 98955), and pQE-GnRH:gD (ATCC Accession

No. 98954). These plasmids are able to express fusion proteins of this invention in *E. coli* bacteria.

In another embodiment, a vector of the subject invention is a plasmid suitable for *in vivo* expression of a fusion protein. Plasmids which are able to transfect eukaryotic cells, and which can be used to construct vectors of the subject invention, can be ascertained by those of ordinary skill in the art. Such plasmids can comprise sequences and encode elements which assist in the *in vivo* expression and processing of the fusion proteins in a vaccinated vertebrate. For example, a plasmid of the present invention can comprise a eukaryotic promoter sequence. As another example, a plasmid of the present invention can comprise a sequence encoding a signal attached to the expressed fusion protein, which signal results in the transportation of the expressed fusion protein to the cell membrane and excretion of the fusion protein from the cell into the vaccinated vertebrate's circulatory system. An example of a plasmid which can be used to construct vectors of the subject invention capable of expressing fusion proteins *in vivo* is pCMV (Clontech, Inc., Palo Alto, CA). Other typical eukaryotic expression plasmids that can be engineered to comprise a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the present invention include an inducible mammalian expression system and the cytomegalovirus promoter-enhancer-based systems (Promega, Madison, WE; Stratagene, La Jolla, CA; Invitrogen). Other plasmids useful for preparing vectors expressing fusion proteins of the subject invention *in vivo* can be ascertained by those of ordinary skill in the art. A preferred example of a plasmid of the subject invention capable of *in vivo* expression of a fusion protein is pCMV-gD:GnRH which has been deposited with the ATCC (ATCC Accession No. 203370).

Vectors of the subject invention also include recombinant viruses which comprise a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the present invention. Such viruses can be prepared according to techniques known in the art. They may, for example, be prepared from bacteriophage, the resulting recombinant bacteriophage being useful for expressing and producing the subject fusion proteins *in vitro* in bacteria. Examples of bacteriophage which can be used to prepare vectors of this invention include T4, T7, ϕ X174, G4, M13, and fd. Other bacteriophage useful for the subject invention may be ascertained by those of ordinary skill in the art.

Recombinant viruses capable of transfecting insect cells or yeast cells can also be constructed for *in vitro* expression and production of fusion proteins of this invention in insect cells and yeast cells, respectively. In this regard, another example of a vector which can be used for *in vitro* production of the fusion proteins of this invention is a recombinant virus based on a baculovirus. In preferred embodiments, the subject invention provides baculovirus vectors which express tmgD-4GnRH, 4GnRH-tmgD-4GnRH, or 4GnRH-tmgD. In a preferred

EXPRESS MAIL NO. EL162820034US

embodiment of this invention, the vector is the baculovirus vector Bac-gD:GnRH, which expresses a tmgD-4GnRH fusion protein. Bac-gD:GnRH has been deposited with the ATCC (ATCC Accession No. VR-2633).

Recombinant viruses capable of infecting and expressing the subject fusion proteins
5 in eukaryotic cells, such as avian or mammalian cells, including viruses for both *in vitro* and *in vivo* expression of the fusion proteins in eukaryotic cells, can also be constructed according to techniques well known in the art. Examples of viruses from which such recombinant viruses can be prepared include poxviruses, such as vaccinia virus, and adenovirus. Both recombinant vaccinia virus and recombinant adenovirus can be used for either *in vitro* or *in vivo* expression. Other viruses suitable for expression in eukaryotic cells can be ascertained
10 by those of ordinary skill in the art.

In another embodiment, a vector of the subject invention is a "transfer vector" comprising a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein of the subject invention. A transfer vector is a plasmid comprising a sequence
15 encoding a peptide, which plasmid can infect a suitable host cell, such as a suitable insect or mammalian cell, in an *in vitro* co-infection process with a virus, causing the host cell to produce a recombinant virus, which recombinant virus is itself a vector that is capable of expressing the peptide encoded by the plasmid in a suitable expression system. Preparation of transfer vectors for *in vitro* production of recombinant virus is well known in the art, and plasmids which are useful for preparing transfer vectors according to this subject invention
20 can be ascertained by those of ordinary skill in the art. Examples of plasmids suitable for preparing transfer vectors include, but are not limited to, pBacPAK8 and pBacPAK 9 (Clontech, Inc.). A preferred transfer vector for preparing a viral vector encoding a fusion protein of the subject invention is the transfer vector pBacHISgD:GnRH.

25 The nucleotide sequence which encodes a fusion protein of the present invention can be ligated to and placed under the control of various nucleotide elements, such as signal sequences, inducible and non-inducible promoters, ribosome binding sites for bacterial expression, and operators. Such elements permit the nucleotide sequence to be transcribed, either *in vivo* or *in vitro*, in a host cell transfected with a vector comprising the polynucleotide
30 molecule, and accordingly to be cloned or expressed in the host cell. Regulatory sequences and enhancer sequences can also be included in the polynucleotide molecules of the invention. The coding sequences are placed in "operative association" with the elements that are included in the polynucleotide molecules, which means that their placement and orientation is such that transcription of the coding sequences can occur. Such placement is
35 within the ordinary skill in the art.

EXPRESS MAIL NO. 8616282003405

Regulatory elements of polynucleotide molecules of the present invention can vary in their strength and specificities. Depending on the host/vector system to be utilized, any of a number of suitable transcription and translation elements can be used. For instance, when cloning in mammalian cell systems, promoters isolated from the genome of mammalian cells, e.g., mouse metallothionein promoter, or from viruses that grow in these cells, vaccinia virus 7.5K promoter or Moloney murine sarcoma virus long terminal repeat, can be used. Promoters obtained by recombinant DNA or synthetic techniques can also be used to provide for transcription of the inserted sequence. In addition, expression from certain promoters can be elevated in the presence of particular inducers, e.g., zinc and cadmium ions for metallothionein promoters. Non-limiting examples of transcriptional regulatory regions or promoters include, for bacteria, the β -gal promoter, the T7 promoter, the T5 promoter, the TAC promoter, λ left and right promoters, trp and lac promoters, trp-lac fusion promoters, etc.; for yeast, glycolytic enzyme promoters, such as ADH-I and -II promoters, GPK promoter, PGI promoter, TRP promoter, etc.; and for mammalian cells, SV40 early and late promoters, adenovirus major late promoters, among others.

Specific initiation signals can also be used for translation of inserted coding sequences. These signals typically include an ATG initiation codon and adjacent sequences. In cases where the polynucleotide molecule of the present invention includes its own initiation codon and adjacent sequences are inserted into the appropriate expression vector, no additional translation control signals may be needed. However, in cases where only a portion of a coding sequence is inserted, exogenous translational control signals, including the ATG initiation codon, may be required. These exogenous translational control signals and initiation codons can be obtained from a variety of sources, both natural and synthetic. Furthermore, the initiation codon must be in phase with the reading frame of the coding regions to ensure in-frame translation of the entire insert.

Vectors of this invention can also include repressor genes and operators, which regulate the transcription of mRNA. Examples of operators which can be included in the subject vectors include the lac operator sequence. Other operators are known in the art, and can be included in the vectors of this invention.

Expression vectors can also contain a polynucleotide molecule of this invention which is further engineered to contain polylinker sequences that encode specific protease cleavage sites so that the expressed fusion protein can be released from expressed vector sequences by treatment with a specific protease. For example, the fusion protein vector can include a nucleotide sequence encoding a thrombin or factor Xa cleavage site, among others.

Expression vectors of the subject invention can also comprise nucleotide sequences that encode a polyaminoacid that can assist in purification of a fusion protein from media

EXPRESS MAIL NO. EL16282003405

following expression. An example of such a nucleotide sequence is a nucleotide sequence encoding a 6XHIS tag, such as the nucleotide sequence set forth in SEQ ID NO: 38.

Transformed Cells for Expressing Fusion Proteins

5 The subject invention also provides transformed cells which comprise a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein as described herein. Cells useful for transformation for this invention include bacterial cells, yeast cells, mammalian cells, insect cells, and plant cells. Transformed cells of this invention can be prepared by transfecting a cell with a vector comprising a polynucleotide molecule
10 comprising a nucleotide sequence encoding the fusion protein as described above.

 Host cells useful in practicing the subject invention can be eukaryotic or prokaryotic. Such transformed host cells include, but are not necessarily limited to, microorganisms, such as bacteria, transformed with a recombinant bacteriophage or plasmid; yeast transformed with a recombinant vector; animal cells, such as mammalian cells, infected with a recombinant
15 virus vector, e.g., adenovirus or vaccinia virus, among others; and insect cells transformed with a recombinant virus vector, e.g. a baculovirus vector.

 For expression and harvesting of fusion proteins *in vitro*, bacterial cells can be used as host cells. For example, a strain of *E. coli* can be used, such as, e.g., the DH5 α strain available from the ATCC, Rockville, MD, USA (ATCC Accession No. 31343) or from
20 Stratagene (La Jolla, CA) or the BL21 strain available from microorganism depositories such as the ATCC. Eukaryotic host cells, including yeast cells and vertebrate cells, e.g., from a mouse, hamster, cow, monkey, or human cell line, among others, can also be utilized effectively. Examples of eukaryotic host cells that can be used to express a fusion protein of the invention include Chinese hamster ovary (CHO) cells (e.g., ATCC Accession No. CCL-
25 61), NIH Swiss mouse embryo cells NIH/3T3 (e.g., ATCC Accession No. CRL-1658), and Madin-Darby bovine kidney (MDBK) cells (ATCC Accession No. CCL-22).

 Other cells that are particularly useful for *in vitro* expression and harvesting of fusion proteins of this invention are cells which possess a system for glycosylation of amino acids of proteins. Some examples of cells that have a glycosylation system are insect cells,
30 mammalian cells and yeast cells. Systems from different cell types can provide different patterns of glycosylation for a fusion protein of the invention.

 The recombinant vector of the invention is preferably transformed or transfected into one or more host cells of a substantially homogeneous culture of cells. The vector can be introduced into host cells in accordance with known techniques, such as, e.g., by protoplast
35 transformation, calcium phosphate precipitation, calcium chloride treatment, microinjection, electroporation, transfection by contact with a recombined virus, liposome-mediated

transfection, DEAE-dextran transfection, transduction, conjugation, or microprojectile bombardment, among others. Selection of transformants can be conducted by standard procedures, such as by selecting for cells expressing a selectable marker, *e.g.*, antibiotic resistance, associated with the recombinant expression vector.

5 Once an expression vector is introduced into the host cell, the integration and maintenance of the polynucleotide sequence encoding a fusion protein of the present invention, either in the host cell genome or episomally, can be confirmed by standard techniques, *e.g.*, by Southern hybridization analysis, restriction enzyme analysis, PCR analysis including reverse transcriptase PCR (rt-PCR), or by immunological assay to detect
10 the expected fusion protein product. Host cells containing a polynucleotide coding sequence and/or expressing a fusion protein of the present invention can be identified by any of at least four general approaches that are well-known in the art, including: (i) DNA-DNA, DNA-RNA, or RNA-antisense RNA hybridization; (ii) detecting the presence of "marker" gene functions; (iii) assessing the level of transcription as measured by the expression of specific mRNA
15 transcripts in the host cell; or (iv) detecting the presence of mature polypeptide product, *e.g.*, by immunoassay, as known in the art.

 Once a polynucleotide sequence encoding a fusion protein of the present invention has been stably introduced into an appropriate cell, the transformed cell can be clonally propagated, and the resulting cells can be grown under conditions conducive to the maximum
20 production of the encoded fusion protein. Such conditions typically include growing transformed cells to high density. Where the expression vector comprises an inducible promoter, appropriate induction conditions such as, *e.g.*, temperature shift, exhaustion of nutrients, addition of gratuitous inducers (*e.g.*, analogs of carbohydrates, such as isopropyl- β -D-thiogalactopyranoside (IPTG)), accumulation of excess metabolic by-products, or the like,
25 are employed as needed to induce expression.

 Where the recombinantly-expressed fusion protein is retained inside the host cells, the cells are harvested and lysed, and the product is purified from the lysate under extraction conditions known in the art to minimize protein degradation such as, *e.g.*, at 4°C, or in the presence of protease inhibitors, or both. Where the recombinantly-expressed fusion protein is
30 secreted from the host cells, the exhausted nutrient medium can simply be collected and the fusion protein isolated therefrom.

 The recombinantly-expressed fusion protein can be purified from cell lysates or culture medium, as necessary, using standard methods, including but not limited to one or more of the following methods: ammonium sulfate precipitation, size fractionation, ion
35 exchange chromatography, HPLC, density centrifugation, and affinity chromatography. The recombinantly-expressed fusion protein can be detected based, *e.g.*, on size, or reactivity with

a fusion-protein-specific antibody, or by the presence of a fusion tag, e.g. a 6XHIS tag. The present invention encompasses recombinantly-expressed fusion protein in an unpurified state, as secreted into the culture fluid or as present in a cell lysate, as well as partially or substantially purified recombinant fusion protein, all being useful for practicing the present invention.

Vaccines, including Dual-Function Vaccines, and Methods using Same

Fusion protein, vectors, and transformed cells of the present invention can be used to prepare dual-function vaccines to induce an immunoinhibitory response in a vertebrate against the peptide to which portion (a) of the subject fusion proteins is analogous, while simultaneously protecting against infection by the pathogen from which portion (b) is derived. Such vaccines are also useful in a vertebrate solely for inhibiting a peptide to which portion (a) is analogous.

Thus, in one aspect, this invention provides a dual-function vaccine which comprises a fusion protein as described above, or a vector or a transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, in an amount effective to inhibit the activity of the peptide from which portion (a) is derived and to protect against infection by the pathogen from which portion (b) is derived in a vertebrate which endogenously synthesizes the peptide and which can be pathogenically infected by the pathogen, along with a carrier acceptable for pharmaceutical or veterinary use.

In a preferred embodiment, the subject invention provides a dual-function vaccine for inhibiting GnRH activity in cattle while simultaneously protecting cattle from BHV-1 infection, which comprises a fusion protein according to the subject invention, or a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, wherein portion (a) of the fusion protein is analogous to all or part of a GnRH peptide and wherein portion (b) is analogous to all or part of a BHV-1 antigen, the fusion protein being present in an amount effective to inhibit GnRH activity in cattle and to also protect cattle from BHV-1 infection, along with a carrier acceptable for veterinary use.

The subject invention also provides a method for inhibiting the activity of an endogenously-synthesized peptide in a vertebrate and for protecting the vertebrate from a pathogenic infection which comprises immunizing the vertebrate with an amount of a dual-function vaccine as described above, which amount is effective to inhibit the activity of the peptide and to protect against infection by the pathogen. In a preferred embodiment, the subject invention provides a method for inhibiting sexual characteristics and for protecting against BHV-1 infection in a cow, which comprises vaccinating the cow with a dual-function vaccine as described above comprising a fusion protein comprising a portion (a) analogous to

all or part of a GnRH peptide and a portion (b) analogous to all or part of a BHV-1 antigen, or vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, in an amount effective to inhibit sexual characteristics and protect against BHV-1 infection.

5 In vaccines which comprise a fusion protein of the invention wherein portion (b) is analogous to all or part of a BHV-1 antigen, the vertebrate which is vaccinated need not be a vertebrate which BHV-1 is capable of pathogenically infecting. In such vertebrates, portion (b) simply acts as a carrier to induce an immune response inhibiting the peptide to which it is connected.

10 Thus, the subject invention also provides a vaccine for inhibiting the activity of a peptide in a vertebrate which comprises a fusion protein of the invention wherein portion (a) is analogous to all or part of a peptide and portion (b) is analogous to all or part of a BHV-1 antigen, or a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, in an amount effective to inhibit the
15 activity of the peptide, along with a carrier acceptable for pharmaceutical or veterinary use.

 In a preferred embodiment, the invention provides a vaccine for inhibiting the activity of GnRH in a vertebrate which comprises a fusion protein wherein portion (a) is analogous to all or part of a GnRH peptide and portion (b) is analogous to all or part of a BHV-1 antigen, or vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide
20 sequence encoding such a fusion protein, in an amount effective to inhibit GnRH activity, along with a carrier acceptable for pharmaceutical or veterinary use.

 The subject invention also provides a method for inhibiting the activity of a peptide, including, but not limited to, the hormone GnRH, in a vertebrate, which comprises immunizing the vertebrate with an amount of the above described vaccine comprising a fusion protein, or
25 a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which fusion protein comprises a proteinaceous portion analogous to all or part of a BHV-1 antigen as a carrier, which amount is effective to inhibit the activity of the peptide.

 The subject invention also provides a method for inhibiting sexual characteristics in a
30 vertebrate, preferably a mammal, which comprises immunizing the vertebrate with an amount of a vaccine comprising a fusion protein comprising a portion (a) analogous to all or part of a GnRH peptide and a portion (b) analogous to all or part of a BHV-1 antigen, or a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which amount is effective to inhibit sexual characteristics.

35 The vertebrate need not be a member of the bovine species, but can be any vertebrate in

EXPRESS MAIL NO. EL16282003405

which GnRH is endogenously synthesized, such as a sheep, pig, horse, goat, dog, cat, or human.

“Sexual characteristics” refers to those characteristics in a vertebrate associated with the vertebrate’s gender and/or the vertebrate’s ability to reproduce, which characteristics are induced, either in whole or in part, either directly or indirectly, by GnRH. Such characteristics are ascertainable by those of ordinary skill in the art. In male cattle, examples of inhibition of such sexual characteristics include repression of aggressive behavior, suppression of testosterone production, reduced libido, regression of the accessory sex glands (including prostates and seminal vesicles), diminution in the testicular volume, and reduction or cessation of spermatogenesis. In female cattle, inhibition of such sexual characteristics include failure to ovulate and infertility, regression of the reproductive tract, and abortion. In one embodiment, GnRH is inhibited in either a male or a female vertebrate such that the sexual characteristics which are inhibited include a functional reproductive system, the present invention thus providing a form of contraception.

The subject invention also provides a method for inhibiting abnormal cell growth in prostate tissue in a male vertebrate, preferably in a mammal, which comprises immunizing the vertebrate with an amount of a vaccine comprising a fusion protein, or a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which fusion protein comprises a portion (a) analogous to all or part of a GnRH peptide and a portion (b) analogous to all or part of a BHV-1 antigen, which amount is effective to inhibit abnormal prostate cell growth.

Vaccines of the present invention can be formulated following accepted convention to include acceptable carriers for animals, including humans, such as standard buffers, stabilizers, diluents, preservatives, and/or solubilizers, and can also be formulated to facilitate sustained release. Diluents include water, saline, dextrose, ethanol, glycerol, and the like. Additives for isotonicity include sodium chloride, dextrose, mannitol, sorbitol, and lactose, among others. Stabilizers include albumin, among others. Other suitable vaccine vehicles and additives, including those that are particularly useful in formulating modified live vaccines, are known or will be apparent to those skilled in the art. See, e.g., Remington’s Pharmaceutical Science, 18th ed., 1990, Mack Publishing, which is incorporated herein by reference.

Vaccines of the present invention can further comprise one or more additional immunomodulatory components such as, e.g., an adjuvant or cytokine, cholera toxin (CT) or heat labile toxin (LT) among others. Non-limiting examples of adjuvants that can be used in the vaccine of the present invention include the RIBI adjuvant system (Ribi Inc., Hamilton, MT), alum, mineral gels such as aluminum hydroxide gel, oil-in-water emulsions, water-in-oil emulsions such as, e.g., Freund’s complete and incomplete adjuvants, Block copolymer

(CytRx, Atlanta GA), QS-21 (Cambridge Biotech Inc., Cambridge MA), SAF-M (Chiron, Emeryville CA), AMPHIGEN® adjuvant, saponin, Quil A or other saponin fraction, monophosphoryl lipid A, and Avridine lipid-amine adjuvant. Non-limiting examples of oil-in-water emulsions useful in the vaccine of the invention include modified SEAM62 and SEAM 1/2 formulations. Modified SEAM62 is an oil-in-water emulsion containing 5% (v/v) squalene (Sigma), 1% (v/v) SPAN® 85 detergent (ICI Surfactants), 0.7% (v/v) TWEEN® 80 detergent (ICI Surfactants), 2.5% (v/v) ethanol, 200 µg/ml Quil A, 100 µg/ml cholesterol, and 0.5% (v/v) lecithin. Modified SEAM 1/2 is an oil-in-water emulsion comprising 5% (v/v) squalene, 1% (v/v) SPAN® 85 detergent, 0.7% (v/v) Tween 80 detergent, 2.5% (v/v) ethanol, 100 µg/ml Quil A, and 50 µg/ml cholesterol. Other immunomodulatory agents that can be included in the vaccine include, e.g., one or more interleukins, interferons, or other known cytokines. Where the vaccine comprises live transformed cells, the adjuvant is preferably selected based on the ability of the resulting vaccine formulation to maintain at least some degree of viability of the live transformed cells.

A vaccine comprising transformed cells of the present invention can be prepared by standard techniques, for example using an aliquot of culture fluid containing said transformed cells, either free in the medium or residing in mammalian host cells, or both, that can be administered directly, or in concentrated form, to the subject. Alternatively, modified live transformed cells can be combined with a carrier acceptable for pharmaceutical or veterinary use, with or without an immunomodulatory agent, selected from those known in the art and appropriate to the chosen route of administration, where at least some degree of viability of the live cells in the vaccine composition is maintained. Such methods are known in the art.

Where a vaccine of this invention comprises live transformed cells, the vaccine can be stored cold or frozen. Where the vaccine composition comprises a fusion protein, vector, or inactivated transformed cells of the present invention, the vaccine may be stored frozen, or in lyophilized form to be rehydrated prior to administration using an appropriate diluent.

Vaccines of the present invention can optionally be formulated for sustained release of the fusion protein. Examples of such sustained release formulations include fusion protein in combination with composites of biocompatible polymers, such as, e.g., poly(lactic acid), poly(lactic-co-glycolic acid), methylcellulose, hyaluronic acid, collagen and the like. The structure, selection and use of degradable polymers in drug delivery vehicles have been reviewed in several publications, including A. Domb *et al.*, 1992, *Polymers for Advanced Technologies* 3: 279-292, which is incorporated herein by reference. Additional guidance in selecting and using polymers in pharmaceutical formulations can be found in the text by M. Chasin and R. Langer (eds), 1990, "Biodegradable Polymers as Drug Delivery Systems" in: Drugs and the Pharmaceutical Sciences, Vol. 45, M. Dekker, NY, which is also incorporated herein by reference. Alternatively, or additionally, the fusion protein, vector, or transformed

cells can be microencapsulated to improve administration and efficacy. Methods for microencapsulating antigens are well-known in the art, and include techniques described, e.g., in U.S. Patent 3,137,631; U.S. Patent 3,959,457; U.S. Patent 4,205,060; U.S. Patent 4,606,940; U.S. Patent 4,744,933; U.S. Patent 5,132,117; and International Patent Publication
5 WO 95/28227, all of which are incorporated herein by reference.

Liposomes can also be used to provide for the sustained release of fusion protein, vector, or transformed cell. Details concerning how to make and use liposomal formulations can be found in, among other places, U.S. Patent 4,016,100; U.S. Patent 4,452,747; U.S. Patent 4,921,706; U.S. Patent 4,927,637; U.S. Patent 4,944,948; U.S. Patent 5,008,050; and
10 U.S. Patent 5,009,956, all of which are incorporated herein by reference.

An effective amount of any of the above-described vaccines can be determined by conventional means, starting with a low dose of fusion protein, vector, or transformed cell and then increasing the dosage while monitoring the effects. An effective amount may be obtained after a single administration of a vaccine or after multiple administrations of a vaccine. Known
15 factors may be taken into consideration when determining an optimal dose per animal. These include the species, size, age and general condition of the animal, the presence of other drugs in the animal, and the like. The actual dosage is preferably chosen after consideration of the results from other animal studies.

One method of detecting whether adequate immune response has been achieved is to
20 determine seroconversion and antibody titer in the animal after vaccination. The timing of vaccination and the number of boosters, if any, will preferably be determined by a qualified scientist or veterinarian based on analysis of all relevant factors, some of which are described above.

The effective dose amount of fusion protein, vector, and transformed cell of the present
25 invention can be determined using known techniques, taking into account factors that can be determined by one of ordinary skill in the art such as the weight of the animal to be vaccinated. The dose amount of fusion protein of the present invention in a vaccine of the present invention preferably ranges from about 1 μ g to about 10 mg, more preferably from about 50 μ g to about 1 mg, and most preferably from about 100 μ g to about 0.5 mg. The dose amount of a vector of the
30 present invention in a vaccine of the present invention preferably ranges from about 50 μ g to about 1 mg. The dose amount of transformed cells of the present invention in a vaccine of the present invention preferably ranges from about 1×10^3 to about 1×10^8 cells/ml, and more preferably from about 1×10^5 to about 1×10^7 cells/ml. A suitable dosage size ranges from about 0.5 ml to about 10 ml, and more preferably from about 1 ml to about 5 ml.

35 Where inhibiting abnormal cell growth in prostate is concerned, an effective amount of any of the above-described vaccines can be determined by conventional means, starting with a

low dose of fusion protein, vector, or transformed cell and then increasing the dosage while monitoring the effects. Known factors can be taken into consideration when determining an optimal dose per animal. Some factors are described above.

"Abnormal cell growth" means cell growth which is independent of normal regulatory mechanisms (e.g., loss of contact inhibition). This includes the abnormal growth of: (1) malignant prostate tumor cells, such as prostate carcinoma cells, (2) benign cells of other proliferative disorders in prostate tissue, and (3) any other unregulated cell growth in prostate tissue associated with GnRH activity. "Inhibiting prostate carcinoma growth" and like phrases as used herein mean slowing, halting, and/or reversing abnormal cell growth in prostate tissue.

The present invention further provides a method of preparing a vaccine comprising a fusion protein as described above, which method comprises combining an effective amount of a fusion protein of the present invention, with a carrier acceptable for pharmaceutical or veterinary use.

Antibodies

The subject invention further provides a method of making polyclonal antibodies directed against a peptide that is endogenously synthesized in a vertebrate which comprises vaccinating such a vertebrate with an antibody-inducing amount of a fusion protein of the present invention, or a vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which fusion protein comprises a portion (a) analogous to all or part of a peptide endogenously synthesized within the vertebrate; obtaining serum containing polyclonal antibodies from the vaccinated vertebrate; and isolating from the serum polyclonal antibodies which bind to the endogenously-synthesized peptide; thereby making polyclonal antibodies directed against the peptide. Methods for obtaining serum from a vaccinated vertebrate and for isolating specific polyclonal antibodies therefrom are known in the art. In a preferred embodiment, the fusion protein comprises a portion (a) analogous to all or part of a GnRH peptide, and the peptide against which polyclonal antibodies are made is GnRH. The subject invention further provides polyclonal antibodies directed against an endogenously-synthesized peptide made according to this method. In a preferred embodiment, the polyclonal antibodies are directed against GnRH.

The subject invention further provides a method of making a monoclonal antibody directed against a peptide that is endogenously synthesized in a vertebrate which comprises vaccinating such a vertebrate with an antibody-inducing amount of a fusion protein of the present invention, or vector or transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding such a fusion protein, which fusion protein

comprises a portion (a) analogous to all or part of a peptide endogenously synthesized within the vertebrate; and isolating a spleen cell from the vaccinated vertebrate which spleen cell excretes a monoclonal antibody that specifically binds to the endogenously-synthesized peptide; thereby making a monoclonal antibody directed against the peptide. In a preferred embodiment, the fusion protein comprises a portion (a) analogous to all or part of a GnRH peptide, and the peptide against which the monoclonal antibody is made is GnRH. The subject invention further provides monoclonal antibodies directed against an endogenously-synthesized peptide made according to this method. In a preferred embodiment, the monoclonal antibodies are directed against GnRH.

Methods for isolating spleen cells from a vaccinated animal which excrete a specific monoclonal antibody for purposes of making a monoclonal antibody are known in the art. Such methods include, but are not limited to, the hybridoma technique originally described by Kohler and Milstein (Nature, 1975, 256: 495-497); the human B-cell hybridoma technique (Kosbor *et al.*, 1983, Immunology Today 4:72; Cote *et al.*, 1983, Proc. Natl. Acad. Sci. USA 80: 2026-2030); and the EBV-hybridoma technique (Cole, *et al.*, 1985, Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). These publications are incorporated herein by reference.

Techniques for the production of monoclonal antibodies and antibody fragments are additionally described, among other places, in Harlow and Lane, 1988, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, and in J. W. Goding, 1986, Monoclonal Antibodies: Principles and Practice, Academic Press, London, which are incorporated herein by reference.

The following examples are provided to merely illustrate aspects of the subject invention. They are not intended, and should not be construed, to limit the invention set forth in the claims and more fully described herein.

Examples

Example 1: Plasmids Expressing gD/GnRH Fusion Proteins

Construction of pQE-tmgD: The plasmid FlgD/Pots207nco(#79) (encoding a full-length gD, hereinafter "gD/Pots") was digested with NcoI/XbaI and the resulting 1.26 kb fragment was cloned into the corresponding sites of pUC21, generating the plasmid pUC-FLgD. The complete sequence of the NcoI/XbaI fragment in the plasmid pUC-FLgD was determined on both DNA strands using Sanger fluorescent dideoxy chain termination sequencing technology. Figure 3 shows the sequence results and characteristics. The nucleotide sequence encoding gD/Pots is included in SEQ ID NO: 16.

DNA alignment between gD/Pots and published BHV-1gD (GenBank Accession No. M59846) shows 94.7% homology with the majority of the mismatches occurring 3' of the transmembrane domain (Figure 4).

5 Amino acid alignment between gD/Pots and M59846 shows four amino acid differences, one of which is located in the signal sequence and the other three in or around the transmembrane domain (Figure 5).

10 The signal sequence for the gD protein was removed in order to facilitate expression of the protein in *E. coli*. The signal sequence removal was carried out by digesting pUC-FLgD plasmid DNA with Nco I/Hind III and filling in the ends with the Klenow fragment of DNA Polymerase I. The resulting DNA fragment was gel-purified and ligated. Colonies were screened for a shift in mobility of a Sph I/Sac I fragment that would indicate deletion of the 50 bp fragment. Two positive clones were selected and sequenced across the Nco I/Hind III deletion region. All clones were shown to have the correct sequence. Clone #1 was designated pUC-MgD was chosen for further manipulation.

15 The mature gD sequence was subcloned into an *E. coli* expression vector, for production of the protein on a large scale basis. To this end, a 1.07 kb Sph I/Sac I fragment from pUC-MgD containing the mature gD sequence (truncated at the 3' end to exclude the transmembrane domain) was subcloned into the corresponding sites of pQE-31 (Qiagen). (pQE-31 uses the phage T5 promoter and two lac operator sequences for greater repression before induction of expression with IPTG. pQE-31 also contains an N-terminal 6XHIS tag fusion for purification purposes.) The resulting clones were screened for the 1.07 kb Sph I/Sac I fragment. One positive clone, designated pQE-tmgD, was selected for preparation of further plasmids, *infra*. pQE-tmgD encodes an N-terminal 6XHIS tag fused to a truncated mature gD (tmgD) sequence, terminated by a vector-encoded stop codon following the Sac I site. The junction regions of the gD sequence and the plasmid backbone were sequenced to verify the integrity of the insert, and were found to be correct. The sequence encoding the tmgD (not including the 6XHIS tag) in pQE-tmgD is set forth in SEQ ID NO: 36. The amino acid sequence of the tmgD encoded by pQE-tmgD (without the 6XHIS tag) is set forth in SEQ ID NO: 35.

30 *Construction of GnRH -tetramer clones:* Twelve different oligonucleotides (sense and complementary (reverse) strands) encoding GnRH (monomers and dimers) having different terminal DNA sequences were prepared. These twelve oligonucleotides are provided in SEQ ID NOS: 1-12.

35 Oligonucleotides 9 and 10 were annealed and cloned into the BamHI/XhoI sites of pBS KS+ (Stratagene), generating p98BS/GnRH. A plasmid encoding a GnRH tetramer was constructed from plasmid p98BS/GnRH by adding annealed oligonucleotides 7 and 8 at the

Sma I/Xho I sites. A reconstruction of the full length tetramer was necessary because sequence analysis of 5 separate clones showed that all had base changes in the synthetic primer region. A clone containing the full length tetramer with the correct sequence was constructed by replacing a 106 bp Eag I fragment from one mutant clone with the corresponding fragment from a clone lacking base changes in this region. One of the resulting reconstructed clones was sequenced and found to have the correct DNA sequence encoding the GnRH tetramer. This clone contained one sequence difference from the predicted nucleotide sequence for the GnRH tetramer construct. The change is an additional G, 3' and outside the GnRH coding region, and, therefore, does not affect the coding region for GnRH. (The additional G was present in the clone used for the reconstruction, and is likely to be due to an error in the synthetic primer sequence.) This clone was designated p9897-R. A portion of p9897-R, including the sequence encoding the GnRH tetramer, is shown in Figure 2. The sequence encoding the GnRH tetramer is set forth in SEQ ID NO: 32. The amino acid sequence of the GnRH tetramer encoded by p9897-R is set forth in SEQ ID NO: 33.

A PCR was employed using primers P14-S1 (SEQ ID NO:42) and P14-A138 (SEQ ID NO: 43) with template DNA from plasmid p9897-R to generate a 138bp fragment containing the GnRH tetramer PCR fragment having a 3' stop codon and synthetic 5' SacI and 3' HindIII ends. The PCR fragment was cloned into the pGEM-T EASY vector (Promega, Madison, Wisconsin), generating p9897 S/d3. The clone was sequenced and found to have the correct sequence. The clone, p9897 S/d3, provides a source for a GnRH tetramer coding sequence with SacI and HindIII ends for future cloning into pQE vectors.

Construction of pQE-gD:GnRH: A 126 bp SacI/HindIII fragment from p9897 S/d3 containing the GnRH tetramer was cloned into the corresponding sites of plasmid pQE-tmgD. Colonies were screened for the presence of the 126 bp SacI/HindIII fragment and a 1165 bp BamHI/HindIII fragment indicating proper orientation of insert. The junction regions adjacent to the cloning sites were analyzed by DNA sequencing and found to be correct. The nucleotide sequence encoding tmgD-4GnRH, including the 6XHIS tag, and plasmid flanking sequences are set forth in SEQ ID NO: 24. The amino acid sequence of the tmgD-4GnRH encoded by pQE-gD:GnRH is set forth in SEQ ID NO: 25. As described above, tmgD-4GnRH is a fusion construct wherein a GnRH tetramer is fused to the carboxy terminus of truncated mature gD.

Construction of pQE-GnRH:gD: The GnRH tetramer coding sequence in p9897-R was cleaved with BamHI/NcoI, the ends blunted by filling in with Klenow, and the 132 bp fragment was gel purified. A mature gD vector fragment (i.e. without the signal sequence) was prepared by cleavage from pUC-FLgD with NcoI/HindIII, blunting the ends by filling in

with Klenow, and gel purifying the 4.4 kb fragment. After ligation with the 132 bp fragment from p9897-R and transformation, clones were screened for the regeneration of the 5' BamH I and Nco I sites resulting from ligation in the correct orientation. Additional screening for the generation of an ~ 400 bp Nde I fragment confirmed the correct structure. The construct was
5 sequenced across the GnRH/gD junctions to confirm the correct sequence. This construct, designated pUC-GnRH:gD, contains a GnRH tetramer sequence fused to the amino terminus of a mature full-length gD sequence in a pUC vector.

An 1161 bp GnRH tetramer/truncated mature gD fusion sequence was obtained by digesting pUC-GnRH:gD with Sph I and Sac I restriction enzymes. This 1161 bp fragment
10 was cloned into the corresponding sites of pQE-31, generating pQE-GnRH:gD. Clones were screened for the 1161bp Sph I/Sac I fragment, and for the correct pattern of Nde I fragments (380 bp, 2.0, 2.2 kb).

The nucleotide sequence encoding 4GnRH-tmgD, including the 6XHIS tag, and plasmid flanking sequences are set forth in SEQ ID NO: 22. The amino acid sequence of the
15 4GnRH-tmgD encoded by pQE-GnRH:gD is set forth in SEQ ID NO: 23.

Construction of pQE-GnRH:gD:GnRH: The 126 bp Sac I/Hind III fragment from p9897 S/d3 was subcloned into the corresponding sites of plasmid pQE-GnRH:gD, generating pQE-GnRH:gD:GnRH. Clones were screened for the 126 bp Sac I/Hind III fragment, as well
20 as for the correct pattern of Nde I fragments.

pQE-GnRH:gD:GnRH encodes a 4GnRH-tmgD-4GnRH fusion protein. As described above, 4GnRH-tmgD-4GnRH comprises a truncated mature gD having a GnRH tetramer fused at both the amino and carboxy termini. The nucleotide coding sequence and flanking sequences from pQE-GnRH:gD:GnRH are provided in SEQ ID NO: 26. The amino acid sequence of the 4GnRH-tmgD-4GnRH encoded by pQE-GnRH:gD:GnRH, including the
25 6XHIS tag is set forth in SEQ ID NO: 27.

Comparison of expression products from bacterial expression vector pQE constructs:

All four constructs contained a tmgD derived from clone FlgD/Pots207nco(#79), which included amino acids 19 through 358 of FlgD/Pots207nco(#79).

All four constructs contained an amino terminal pQE-HIS leader sequence (a 6XHIS tag) denoted by amino acid designation: MRGSHHHHHHTDPHA (SEQ ID NO: 37). The
30 coding sequence for the 6XHIS tag is set forth in SEQ ID NO:38.

All four constructs had a 2 or 3 amino acid linker after the 6XHIS leader sequence.

All GnRH products were derived from GnRH tetramer clone p9897-R.

The pQE-GnRH:gD and pQE-GnRH:gD:GnRH clones contained a three amino acid
35 linker (SMS) between the amino terminal GnRH tetramer and the tmgD sequence.

The pQE-gD and pQE-GnRH:gD clones contained an extra ten amino acids at the carboxy terminal end of tmgD from the vector sequence as an artifact from cloning.

The pQE-gD:GnRH and pQE-GnRH:gD:GnRH clones contained a one amino acid (proline) linker between tmgD and GnRH carboxy fusion.

5 See Figure 10 for an illustration of each of the pQE constructs.

Example 2: Expression of GnRH/gD Fusion Proteins by Transformed Bacterial Cells

All of the pQE constructs described in Example 1, above, were transformed into E.coli DH5 α -F'IQ cells for expression. For induction of expression, cells were grown to an
10 OD₆₀₀ of 0.7-0.9 in a 2 liter baffled culture flask in 2xYT broth containing 100 μ g/ml Ampicillin and 25 μ g/ml Kanamycin, then induced with 1-2mM IPTG and incubated for 4 hours at 37 degrees Celsius. Average OD₆₀₀ readings at harvest time were 1.3. Expression of all four constructs was confirmed by Western blot analysis.

Example 3: Formulation of Fusion Protein Vaccines and Immunization of Mice

Vaccine Assembly: Fusion proteins from pQE-tmgD (as a control), pQE-GnRH:gD, pQE-GnRH:gD:GnRH, and pQE-gD:GnRH were concentrated from inclusion body preparations by preparative electrophoresis on 9% polyacrylamide gels. Bands cut from SDS
20 PAGE gels were dissolved in 25mM Tris, pH 8.3, 192mM glycine and 0.1% SDS (w/v). The equivalent of 10 μ g gD/mouse dose was adjuvanted with SEAM1 (Squalene Emulsion Adjuvant Metabolizable) emulsion (10 μ g QuilA/100 μ l dose). Vaccine formulations were stored at 4°C. SEAM1 is 5% squalene, 0.1% Vitamin E acetate, 1% Span 85, 0.70% Tween 80, 2mg/ml QuilA, and 400 μ l/ml cholesterol.

25 *Mice:* BALB/c males were used in the study after they were 8 weeks of age (10/group). Mice were initially housed in groups of 10, however, controls were subsequently moved to individual cages to prevent fighting.

Immunization: Mice were immunized subcutaneously with 10 μ g fusion protein in 100 μ L adjuvant, described above. Three immunizations were given at study days 0, 20, and
30 41.

Anti-GnRH antibodies by ELISA: Serum samples were collected at study days 0, 20, 31, 41, 55, 62, 69, and 146 and were evaluated for anti GnRH antibody titers in a peptide ELISA (enzyme linked immunoabsorbant assay). A biotinylated GnRH peptide (Biotin-GnRH) (0.1 μ g/mL in 25 mM Tris, 0.15 M NaCl at pH 7.6) consisting of the natural sequence plus a 4
35 amino acid linker (CAGAEHWSYGLRPG), purified by HPLC on a reverse phase column, was adsorbed to avidin coated plates and incubated at room temperature for 2 hours. Excess

peptide was removed by washing plates four times with the wash buffer (25mM Tris, 0.15M NaCl, 0.05% Tween-20 and 0.05% BSA (bovine serum albumin) fraction V). Then, five-fold serial dilutions of positive control, negative and unknown mouse sera in diluent (25 mM Tris, 0.15 M NaCl, .05% BSA) (100 µl /well) were added to the peptide coated wells and incubated for 30 minutes at room temperature. Plates were washed four times in wash buffer and then rabbit anti mouse IgG (IgG specific)-horseradish peroxidase (Zymed, California) was added to each well (1:4,000, 100 µl/well). After incubation for 30 minutes at room temperature the bound antibody was detected with 3,3',5,5'-tetramethyl benzidine substrate (Kierkegaard & Perry, cat#50-76-04) (100 µl/well, 15 minutes in the dark) and the reaction was halted with the addition of 50 µl/well of 0.18 M H₂SO₄. Absorbance at 450 nm was measured with a Molecular Devices microplate reader. To calculate antibody titers, a positive control curve is generated and titers of unknown samples are extrapolated from the curve using computer software.

BHV-1 gD ELISA: Serum samples were collected at study days 0, 20, 31, 41, 55, 62, 69, and 146 and were evaluated for anti gD BHV-1 antibody titers by ELISA. Purified recombinant gD BHV-1 expressed from MDBK (Madin Darby Bovine Kidney) cells (1 µg/mL in Dulbecco's PBS + 0.01% thimerosal, 100µL/well) was adsorbed onto microtiter plates for 18-24 hours at 4°C. Excess protein was washed from wells then unbound sites in wells were blocked by incubating for 2 hours at 37°C with 300µl of 1%PVA (polyvinyl acetate) in DPBS (Dulbecco's phosphate buffered saline) with 0.01% thimerosal. Serum samples (positive and negative control and unknown serum) were diluted 1:50, then serially by 4-fold dilutions in 1%PVA in DPBS with 0.01% thimerosal and 100µl added to each well. The assay was incubated 45 minutes at 37°C. Plates were washed four times with distilled H₂O, then HRP (horse radish peroxidase) goat anti-mouse (1:10000 in 1%PVA in DPBS with 0.01% thimerosal, 100µl/well, KP+L) was added and plates were incubated 30 minutes at 37°C. Wells were washed four times with distilled H₂O then the assay was developed with ABTS (2,2'-azino-di[3-ethyl-benzthiazoline sulfonate (6) substrate (100µl/well, RT, 15min). The reaction was read at 405/490nm on an ELISA reader. Titers were calculated using the Forecast method in EXCEL™ (Microsoft, Redmond, Washington) using 0.5 OD as a cutoff and using 2 dilutions above 0.5 and 1 dilution below the 0.5 OD to extrapolate titers.

Testosterone Concentrations: Serum samples from study days 0, 41 and 69 were evaluated for testosterone concentrations. The assay was a human testosterone radioimmunoassay using antibody that cross-reacts with murine testosterone. Human testosterone standards are used in the assay. The murine samples tend to run at the lower

end of the human testosterone standard curve, leading to a wider variability in normal values. The sensitivity of the assay is 0.02ng/mL.

Necropsy and Histopathology: Animals were sacrificed at study day 146. Testes, epididymides and prostate with seminal vesicle were removed and weighed prior to fixation of tissues in Bouin's fixative [75mL picric acid (saturated solution); 25mL formalin (37%); 5mL acetic acid (4.76%)]. Tissues were fixed for 48 hours then rinsed in 50% ethanol:H₂O. Tissues were stored in fresh 50% ethanol prior to analysis. Tissues were processed and embedded in paraffin and 5µm sections cut and stained with hematoxylin and eosin. Each organ was evaluated for inflammation, atrophy, and spermatogonial degeneration. Scores were assigned based on the level of aspermatogenesis, atrophy, or other lesions. Weights were scored as a percentage of the mean weight in the normal control group. A cumulative score was assigned to each animal.

Results:

Anti-gD antibody responses: All mice that were immunized with gD or a gD-containing fusion protein generated anti-gD ELISA antibodies, regardless of whether gD was expressed in procaryotic (i.e. E.coli expressed carboxyl, amino or carboxyl-amino fusion protein) or eucaryotic expression systems (i.e. MDBK expressed protein).

Anti-GnRH antibody responses: A hierarchy of anti-GnRH titers were induced by the different fusion proteins: tmgD-4GnRH (i.e. having a GnRH tetramer at the carboxy end of the protein) generated the highest titers followed by 4GnRH-tmgD-4GnRH, while the lowest titers were induced in the 4GnRH-tmgD immunized. In all groups anti-GnRH titers peaked after the second immunization and remained at plateau for greater than 2 months.

All (9 of 9) mice immunized with the tmgD-4GnRH made antibody responses to GnRH when measured by peptide ELISA, although 2/9 mice were low responders. There were 3/10 nonresponders in the 4GnRH-tmgD group and 1/9 nonresponders in the 4GnRH-tmgD-4GnRH. All the GnRH nonresponders were gD responders.

Effect of anti-GnRH antibodies on the male reproductive system: To determine whether induction of anti-GnRH antibodies would abrogate GnRH function we evaluated testosterone levels before and after GnRH immunization. At necropsy, reproductive tract tissues were weighed then submitted for gross and histological examination. The normal ranges of testosterone concentrations in mice varied widely as measured using the human testosterone radioimmunoassay. However, mice immunized with tmgD-4GnRH had significantly lower mean testosterone concentrations when compared to normal controls or other treatment groups. The prostate, testes and epididymides of tmgD-4GnRH immunized mice were significantly atrophied when gross tissue weight and histological examination of

sperm development was evaluated. Mice immunized with 4GnRH-tmgD-4GnRH were less affected when compared to normal controls.

Example 4: Baculovirus constructs encoding gD/GnRH Fusion Proteins

5 *Construction of pBachISgD:LH and bac-gD:GnRH:* pQE-gD:GnRH (see Example 1) was digested with Hind III, the site blunt-ended by Klenow treatment, and subsequently digested with EcoRI. An approximate 1.2 kb fragment which contained the tmgD-4GnRH coding sequence was gel purified and cloned into STUI/EcoRI digested transfer vector pBacPAK9 plasmid (Clontech, Inc.), forming pBachISgD:LH. (The transfer vector contains
10 sequences compensating for replication deficiency in a replication deficient baculovirus.)

Sf21 insect cells were co-transfected with pBachISgD:LH and replication deficient baculovirus viral DNA. These transfected Sf21 cells generate a recombinant baculovirus (designated bac-gD:GnRH) (ATCC Accession No. VR-2633), which encodes a tmgD-4GnRH fusion protein. Recombination (exchange of DNA) between the transfer vector
15 pBachISgD:LH and replication deficient baculovirus viral DNA is mediated by homologous flanking viral sequences present in pBacPAK9 which allows for efficient transfer of the entire expression cassette (sequence encoding tmgD-4GnRH) from pBachISgD:LH into viral DNA along with the gene or genes that complements for replication deficiency.

Recombinant virus can be purified by plaque assay from infected Sf21 cells.
20 Repeated cycles of Sf21 cell infection and plaque assay purification can be performed to obtain greater concentration of recombinant virus expressing fusion protein for large scale production of the fusion protein. Expression of the recombinant constructs was confirmed by Western blot. Infected Sf21 cells can be collected by centrifugation and transferred to -80° Celsius until processed for recombinant baculovirus.

25 The nucleotide sequence encoding the ORF for the 6XHIS tag, truncated mature gD and GnRH tetramer in bac-gD:GnRH is set forth in SEQ ID NO: 39. Nucleotides #1-45 encode a 6XHIS tag, nucleotides #46-1074 encode a truncated mature BHV-1 gD, nucleotides #1075-1194 encode a GnRH tetramer, and nucleotides #1195-1197 are a stop codon. The amino acid sequence of the fusion protein encoded by bac-gD:GnRH is the same as the
30 sequence set forth in SEQ ID NO: 25.

Construction of pBachISMgD: A recombinant baculovirus construct containing gD was generated as a control. Plasmid pCMV-MgD (see Example 5, *infra*) was digested with PacI and Apal allowing for the isolation of a 950bp fragment containing the majority of the gD gene minus the 5' end. Plasmid pBachISgD:LH underwent digestion with PacI and Apal
35 allowing for the isolation of a 5.6kb fragment containing the plasmid backbone and the 5'

portion of gD. Ligation of the 5.6kb fragment with the 950bp fragment generated plasmid pBachISMgD containing truncated mature gD in transfer vector, pBacPAC9.

Sf21 cells were co-transfected with pBachISMgD and replication deficient virus. These transformed Sf21 cells generate recombinant baculovirus (designated Bac-MgD) which
5 encodes tmgD. Recombinant virus was purified and stored as described above.

Expression: Recombinant baculovirus can be obtained from lysates of infected Sf21 cells. The lysate also contains the fusion protein expressed by the recombinant virus, and the fusion protein may be purified from the lysate. For example, after detergent lysis of the cell pellet, the lysate pellet in the aforementioned example was solubilized in 8 M urea, 50mM
10 Tris, pH 7.5 and loaded onto a Ni NTA column; the tmgD-4GnRH was eluted in a pH step gradient. The lysate, containing both the recombinant baculovirus and fusion protein, can be stored, for example, at -80° Celsius.

Example 5: Plasmid suitable for *in vivo* expression of gD/GnRH Fusion Proteins

The β -Gal gene from pCMV β vector (Clontech, Inc) was removed by EcoRV/NotI
15 restriction digest and the resulting NotI vector fragment was isolated by gel electrophoresis. A synthetic linker containing multiple cloning sites (MC) with NotI ends was cloned into this NotI vector fragment creating pCMV-MC.

A truncated gD gene including the signal sequence was PCR amplified from FlgD/Pots207nco(#79) using primers that introduced an EcoRV site at 5' end, a second codon
20 repaired to encode Gln rather than Glu, a stop codon added after Pro 337 of the coding sequence, and a KpnI site added at the 3' end. This 1083 bp PCR fragment was cloned into EcoRV/KpnI digested pGEM-T EASY vector (Promega, Madison, Wisconsin), generating pGEM-T-EASY/gD, and subsequently sequenced by fluorescent di-deoxy termination chemistry in both directions to ensure integrity of PCR product. The truncated gD fragment
25 was isolated from the pGEM-T-EASY/gD clone by EcoRV/KpnI digestion and subcloned into pCMV-MC. The resulting clone, designated pCMV-gD, was verified by restriction enzyme analysis.

To construct pCMV-gD:GnRH (ATCC accession No. 203370), pQE-gD:GnRH was cleaved with HindIII, blunt ended with Klenow and then digested with Apal. The resulting
30 blunt-ended/Apal 1.05 kb fragment containing tmgD and GnRH tetramer was isolated. Clone, pCMV-gD was cleaved with SmaI, followed by Apal, removing the truncated gD encoding region. The remaining 3.7 kb pCMV vector fragment containing the signal sequence for gD was isolated and used in a ligation reaction with the 1.05 kb fragment containing tmgD and GnRH tetramer. The resulting clone was designated pCMV-gD:GnRH. The ORF encoding
35 the tgD-4GnRH, including the signal sequence, from pCMV-gD:GnRH is set forth in SEQ ID

EXPRESS MAIL NO. 2L162 82003405

NO: 28. The amino acid sequence of the tgD-4GnRH, including the signal sequence, encoded by pCMV-gD:GnRH is set forth in SEQ ID NO: 29.

All patents, patent applications, and publications cited above are incorporated herein by reference in their entirety.

5 The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing
10 description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

DEPOSIT OF BIOLOGICAL MATERIALS

15 The following biological material was deposited with the American Type Culture Collection (ATCC) at 10801 University Blvd., Manassas, Virginia, 20110-2209, USA, on October 22, 1998 and were assigned the following accession numbers:

	<u>Plasmid</u>	<u>Accession No.</u>
	plasmid pQE-gD:GnRH	98953
	plasmid pCMV-gD:GnRH	203370
	plasmid pQE-GnRH:gD	98954
20	plasmid pQE-GnRH:gD:GnRH	98955
	<u>Vector</u>	<u>Accession No.</u>
	baculovirus bac-gD:GnRH	VR-2633

25 All patents, patent applications, and publications cited above are incorporated herein by reference in their entirety.

30 The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

EXPRESS MAIL NO. EL16282 0034 US

What is claimed is:

1. A fusion protein for producing a dual immune response in a vertebrate, which fusion protein comprises:

(a) a first proteinaceous portion analogous to all or part of a peptide endogenously synthesized within the vertebrate, the activity of which peptide is to be inhibited within the vertebrate, and which proteinaceous portion by itself is incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to

(b) a second proteinaceous portion analogous to all or part of an immunogen from a pathogen, which pathogen is capable of pathogenically infecting the vertebrate;

the portion (b) causing the vertebrate's immune system to recognize the portion (a) and produce a response that:

(i) inhibits the activity of the peptide endogenously synthesized within the vertebrate; and

(ii) protects the vertebrate from infection by the pathogen, when the vertebrate is vaccinated with an effective amount of the fusion protein.

2. A fusion protein according to claim 1 comprising a portion (a) analogous to all or part of a GnRH peptide and a portion (b) analogous to all or part of a BHV-1 antigen.

3. A fusion protein for producing an immune response in a vertebrate, which fusion protein comprises:

(a) a first proteinaceous portion analogous to all or part of a peptide the activity of which is to be inhibited within the vertebrate, and which proteinaceous portion by itself is incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to

(b) a second proteinaceous portion analogous to all or part of a BHV-1 antigen;

the second proteinaceous portion (b) causing the vertebrate's immune system to recognize the first proteinaceous portion (a) and produce an immune response capable of inhibiting the activity of the peptide within the vertebrate when the vertebrate is vaccinated with an effective amount of the fusion protein.

4. A fusion protein according to claim 3 comprising a portion (a) analogous to all or part of a GnRH peptide.

5. A fusion protein according to claim 3 wherein portion (b) is analogous to all or part of BHV-1 gD.

6. A polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein according to claim 1 or 3.

EXPRESS MAIL NO. EL162820034US

7. A vector comprising a polynucleotide molecule according to claim 6.
8. A vector according to claim 7, suitable for *in vitro* expression of the fusion protein.
9. A vector according to claim 7, suitable for *in vivo* expression of the fusion protein.
10. A transformed cell comprising a polynucleotide molecule comprising a nucleotide sequence encoding a fusion protein according to claim 1 or 4.
11. A dual-function vaccine which comprises a fusion protein according to claim 1, a vector according to claim 7, or a transformed cell according to claim 10 in an amount effective to inhibit the activity of the peptide from which portion (a) of the fusion protein is derived and to protect against infection by the pathogen from which portion (b) of the fusion protein is derived in a vertebrate which endogenously synthesizes the peptide and which can be pathogenically infected by the pathogen, along with a carrier acceptable for pharmaceutical or veterinary use.
12. A dual-function vaccine for inhibiting GnRH activity in cattle and for protecting cattle against BHV-1 infection which comprises a fusion protein according to claim 2, a vector according to claim 7, or a transformed cell according to claim 10 in an amount effective to inhibit GnRH activity and protect cattle against BHV-1 infection, along with a carrier acceptable for pharmaceutical or veterinary use.
13. A vaccine for inhibiting the activity of a peptide in a vertebrate which comprises a fusion protein according to claim 3, a vector according to claim 7, or a transformed cell according to claim 10 in an amount effective to inhibit the activity of the peptide, along with a carrier acceptable for pharmaceutical or veterinary use.
14. A method for inhibiting the activity of an endogenously-synthesized peptide in a vertebrate and for protecting the vertebrate from a pathogenic infection which comprises immunizing the vertebrate with an amount of a vaccine according to claim 11, which amount is effective to inhibit the activity of the peptide and to protect against infection by the pathogen.
15. A method for inhibiting sexual characteristics in a cow and for protecting the cow against BHV-1 infection which comprises immunizing the cow with an amount of a vaccine according to claim 12, which amount is effective to inhibit sexual characteristics and protect against BHV-1 infection.
16. A method for inhibiting the activity of a peptide in a vertebrate which comprises immunizing the vertebrate with an amount of a vaccine according to claim 13, which amount is effective to inhibit the activity of the peptide.
17. A method for inhibiting sexual characteristics in a vertebrate which comprises immunizing the vertebrate with an amount of a vaccine according to claim 13, wherein the

Year	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	

EXPRESS MAIL NO. EL162820034US

**FUSION PROTEINS COMPRISING CARRIERS THAT CAN
INDUCE A DUAL IMMUNE RESPONSE**


5

Abstract

10 The subject invention provides a fusion protein for producing a dual immune response
in a vertebrate, which fusion protein comprises: (a) a first proteinaceous portion analogous to
all or part of a peptide endogenously synthesized within the vertebrate, the activity of which
peptide is to be inhibited within the vertebrate, and which proteinaceous portion by itself is
incapable of eliciting an effective immunoinhibitory response in said vertebrate; connected to
15 (b) a second proteinaceous portion analogous to all or part of an immunogen from a
pathogen, which pathogen is capable of pathogenically infecting the vertebrate; the portion (b)
causing the vertebrate's immune system to recognize the portion (a) and produce a response
that: (i) inhibits the activity of the peptide endogenously synthesized within the vertebrate; and
20 (ii) protects the vertebrate from infection by the pathogen, when the vertebrate is vaccinated
with an effective amount of the fusion protein. The subject invention also provides fusion
proteins which comprise a proteinaceous portion (b) that is a carrier that is analogous to all or
part of a BHV-1 antigen, which fusion proteins induce in a vertebrate vaccinated with an
effective amount of such fusion protein an immune response that inhibits the activity of a
peptide as recited in (a), above.

EXPRESS MAIL NO. EL162820034US

FIG. 1



+




FIG. 2

+

FIG. 3A




FIG. 3B




FIG. 4A

	10	20	30	40
1	ATGGAAGGGCCGACATTGGCCGTGCTGGGCGCGCTGCTCG			
1	ATGCAAGGGCCGACATTGGCCGTGCTGGGCGCGCTGCTCG			
	50	60	70	80
41	CCGTTGCGGTAGAGCTTGCCTACACCCGCGCCGCGGGTGAC			
41	CCGTTGCGGTGAGCTTGCCTACACCCGCGCCGCGGGTGAC			
	90	100	110	120
81	GGTATACGTGACCCGCGGCGGTACCCGATGCCGCGATAC			
81	GGTATACGTGACCCGCGGCGGTACCCGATGCCGCGATAC			
	130	140	150	160
121	AACTACACTGAACGCTGGCACACTACCGGGGCCCATACCGT			
121	AACTACACTGAACGCTGGCACACTACCGGGGCCCATACCGT			
	170	180	190	200
161	CGCCCTTCGCAGACGGCCGCGAGCAGCCCGTCGAGGTGCG			
161	CGCCCTTCGCAGACGGCCGCGAGCAGCCCGTCGAGGTGCG			
	210	220	230	240
201	CTACGCGACGAGCGCGGCGGCGTGCGACATGCTGGGCGCTG			
201	CTACGCGACGAGCGCGGCGGCGTGCGACATGCTGGGCGCTG			
	250	260	270	280
241	ATCGCAGACCCGCGAGGTGGGGCGCACGCTGTGGGAAGCGG			
241	ATCGCAGACCCGCGAGGTGGGGCGCACGCTGTGGGAAGCGG			
	290	300	310	320
281	TACGCCGGCACGCGCGCGCGGTACAACGCCACGGTCATATG			
281	TACGCCGGCACGCGCGCGCGGTACAACGCCACGGTCATATG			
	330	340	350	360
321	GTACAAGATCGAGAGCGGGTGCGCCCGGCCGCTGTACTAC			
321	GTACAAGATCGAGAGCGGGTGCGCCCGGCCGCTGTACTAC			
	370	380	390	400
361	ATGGAGTACACCGAGTGCGAGCCCAGGAAGCACTTTGGGT			
361	ATGGAGTACACCGAGTGCGAGCCCAGGAAGCACTTTGGGT			
	410	420	430	440
401	ACTGCCGCTACCGCACACCCCCGTTTTGGGACAGCTTCCT			
401	ACTGCCGCTACCGCACACCCCCGTTTTGGGACAGCTTCCT			

12345678910111213141516171819202122232425262728293031323334353637383940414243444546474849505152535455565758596061626364656667686970717273747576777879808182838485868788899091929394959697989910010110210310410510610710810911011111211311411511611711811912012112212312412512612712812913013113213313413513613713813914014114214314414514614714814915015115215315415515615715815916016116216316416516616716816917017117217317417517617717817918018118218318418518618718818919019119219319419519619719819920020120220320420520620720820921021121221321421521621721821922022122222322422522622722822923023123223323423523623723823924024124224324424524624724824925025125225325425525625725825926026126226326426526626726826927027127227327427527627727827928028128228328428528628728828929029129229329429529629729829930030130230330430530630730830931031131231331431531631731831932032132232332432532632732832933033133233333433533633733833934034134234334434534634734834935035135235335435535635735835936036136236336436536636736836937037137237337437537637737837938038138238338438538638738838939039139239339439539639739839940040140240340440540640740840941041141241341441541641741841942042142242342442542642742842943043143243343443543643743843944044144244344444544644744844945045145245345445545645745845946046146246346446546646746846947047147247347447547647747847948048148248348448548648748848949049149249349449549649749849950050150250350450550650750850951051151251351451551651751851952052152252352452552652752852953053153253353453553653753853954054154254354454554654754854955055155255355455555655755855956056156256356456556656756856957057157257357457557657757857958058158258358458558658758858959059159259359459559659759859960060160260360460560660760860961061161261361461561661761861962062162262362462562662762862963063163263363463563663763863964064164264364464564664764864965065165265365465565665765865966066166266366466566666766866967067167267367467567667767867968068168268368468568668768868969069169269369469569669769869970070170270370470570670770870971071171271371471571671771871972072172272372472572672772872973073173273373473573673773873974074174274374474574674774874975075175275375475575675775875976076176276376476576676776876977077177277377477577677777877978078178278378478578678778878979079179279379479579679779879980080180280380480580680780880981081181281381481581681781881982082182282382482582682782882983083183283383483583683783883984084184284384484584684784884985085185285385485585685785885986086186286386486586686786886987087187287387487587687787887988088188288388488588688788888989089189289389489589689789889990090190290390490590690790890991091191291391491591691791891992092192292392492592692792892993093193293393493593693793893994094194294394494594694794894995095195295395495595695795895996096196296396496596696796896997097197297397497597697797897998098198298398498598698798898999099199299399499599699799899910001001100210031004100510061007100810091010101110121013101410151016101710181019102010211022102310241025102610271028102910301031103210331034103510361037103810391040104110421043104410451046104710481049105010511052105310541055105610571058105910601061106210631064106510661067106810691070107110721073107410751076107710781079108010811082108310841085108610871088108910901091109210931094109510961097109810991100110111021103110411051106110711081109111011111112111311141115111611171118111911201121112211231124112511261127112811291130113111321133113411351136113711381139114011411142114311441145114611471148114911501151115211531154115511561157115811591160116111621163116411651166116711681169117011711172117311741175117611771178117911801181118211831184118511861187118811891190119111921193119411951196119711981199120012011202120312041205120612071208120912101211121212131214121512161217121812191220122112221223122412251226122712281229123012311232123312341235123612371238123912401241124212431244124512461247124812491250125112521253125412551256125712581259126012611262126312641265126612671268126912701271127212731274127512761277127812791280128112821283128412851286128712881289129012911292129312941295129612971298129913

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)	(25)	(26)	(27)	(28)	(29)	(30)	(31)	(32)	(33)	(34)	(35)	(36)	(37)	(38)	(39)	(40)	(41)	(42)	(43)	(44)	(45)	(46)	(47)	(48)	(49)	(50)	(51)	(52)	(53)	(54)	(55)	(56)	(57)	(58)	(59)	(60)	(61)	(62)	(63)	(64)	(65)	(66)	(67)	(68)	(69)	(70)	(71)	(72)	(73)	(74)	(75)	(76)	(77)	(78)	(79)	(80)	(81)	(82)	(83)	(84)	(85)	(86)	(87)	(88)	(89)	(90)	(91)	(92)	(93)	(94)	(95)	(96)	(97)	(98)	(99)	(100)
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-------

FIG. 5

	10	20	30	40																																				
1	M	C	G	P	T	L	A	V	L	G	A	L	L	A	V	A	V	S	L	P	T	P	A	P	R	V	T	V	Y	V	D	P	P	A	Y	P	M	P	R	Y
1	M	E	G	P	T	L	A	V	L	G	A	L	L	A	V	A	V	S	L	P	T	P	A	P	R	V	T	V	Y	V	D	P	P	A	Y	P	M	P	R	Y
	50	60	70	80																																				
41	N	Y	T	E	R	W	H	T	T	G	P	I	P	S	P	F	A	D	G	R	E	Q	P	V	E	V	R	Y	A	T	S	A	A	A	C	D	M	L	A	L
41	N	Y	T	E	R	W	H	T	T	G	P	I	P	S	P	F	A	D	G	R	E	Q	P	V	E	V	R	Y	A	T	S	A	A	A	C	D	M	L	A	L
	90	100	110	120																																				
81	I	A	D	P	Q	V	G	R	T	L	W	E	A	V	R	R	H	A	R	A	Y	N	A	T	V	I	W	Y	K	I	E	S	G	C	A	R	P	L	Y	Y
81	I	A	D	P	Q	V	G	R	T	L	W	E	A	V	R	R	H	A	R	A	Y	N	A	T	V	I	W	Y	K	I	E	S	G	C	A	R	P	L	Y	Y
	130	140	150	160																																				
121	M	E	Y	T	E	C	E	P	R	K	H	F	G	Y	C	R	Y	R	T	P	P	F	W	D	S	F	L	A	G	F	A	Y	P	T	D	D	E	L	G	L
121	M	E	Y	T	E	C	E	P	R	K	H	F	G	Y	C	R	Y	R	T	P	P	F	W	D	S	F	L	A	G	F	A	Y	P	T	D	D	E	L	G	L
	170	180	190	200																																				
161	I	X	A	A	P	A	R	L	V	E	G	Q	Y	R	R	A	L	Y	I	D	G	T	V	A	Y	T	D	F	M	V	S	L	P	A	G	D	C	W	F	S
161	I	X	A	A	P	A	R	L	V	E	G	Q	Y	R	R	A	L	Y	I	D	G	T	V	A	Y	T	D	F	M	V	S	L	P	A	F	D	C	W	F	S
	210	220	230	240																																				
201	K	L	G	A	A	R	G	Y	T	F	G	A	C	F	P	A	R	D	Y	E	Q	K	K	V	L	R	L	T	Y	L	T	Q	Y	P	Q	E	A	H	K	
201	K	L	G	A	A	R	G	Y	T	F	G	A	C	F	P	A	R	D	Y	E	Q	K	K	V	L	R	L	T	Y	L	T	Q	Y	P	Q	E	A	H	K	
	250	260	270	280																																				
241	A	I	V	D	Y	W	F	M	R	H	G	G	V	V	P	P	Y	F	E	E	S	K	G	Y	E	P	P	P	A	A	D	G	G	S	P	A	P	P	G	D
241	A	I	V	D	Y	W	F	M	R	H	G	G	V	V	P	P	Y	F	E	E	S	K	G	Y	E	P	P	P	A	A	D	G	G	S	P	A	P	P	G	D
	290	300	310	320																																				
281	D	E	A	R	E	D	E	G	E	T	E	D	G	A	A	G	R	E	G	N	G	G	P	P	G	P	E	G	D	G	E	S	Q	T	P	E	A	N	G	G
281	D	E	A	R	E	D	E	G	E	T	E	D	G	A	A	G	R	E	G	N	G	G	P	P	G	P	E	G	D	G	E	S	Q	T	P	E	A	N	G	G
	330	340	350	360																																				
321	A	E	G	E	P	K	P	G	P	S	P	D	A	D	R	P	E	G	W	P	S	L	E	A	I	T	H	P	P	P	A	P	A	T	P	A	-	A	P	D
321	A	E	G	E	P	K	P	G	P	S	P	D	A	D	R	P	E	G	W	P	S	L	E	A	I	T	H	P	P	P	A	P	A	T	P	A	R	A	P	D
	370	380	390	400																																				
360	A	V	P	V	S	V	G	I	G	I	A	A	A	I	A	C	V	A	A	A	A	A	G	A	Y	F	V	Y	T	R	R	R	G	A	G	P	L	P	R	
361	A	V	S	V	S	V	G	I	G	I	A	A	A	I	A	C	V	A	A	A	A	A	G	A	Y	F	V	Y	T	R	R	R	G	A	G	P	L	P	R	
	410																																							
400	K	F	K	K	L	P	A	F	G	N	V	N	Y	S	A	L	P	G	*																					
401	K	F	K	K	L	P	A	F	G	N	V	N	Y	S	A	L	P	G	*																					

GenBank: 64030360

FIG. 6A




FIG. 6B




FIG. 6C

CGCCTACCCACGACGAGCTGGGACTGATTATGCGGCGCCCGCGGCTCGTCGAGGGCCAGTACCGACGCGGC 640
GCGGATGGGGTGCCCTGCTCGACCCCTGACTAATACCGCGCGGGCGCCGAGCAGCTCCCGGTCATGGCTGCGCGCGC

gD coding sequence

Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile Met Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala

6x his:MgD coding seq

TGTACATCGACGGCACGGTCGCCCTATACAGATTTTCATGGTTTCGCTGCCGGGACTGCTGGTTCTCGAAACTCGGC 720
ACATGTAGCTGCCGTGCCAGCGGATATGTCTAAAGTACCAAAAGCAGCGCGGCCCTGACGACCAAGAGCTTTGAGCCG

gD coding sequence

Leu Tyr Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro Ala Gly Asp Cys Trp Phe Ser Lys Leu Gly

6x his:MgD coding seq

GCGGCTCGCGGTACACCTTTGGCGCGTGCTTCCCGGCCGGGATTACGAGCAAAAGAGTTCTGCGCCTGACGTATCT 800
CGCCGAGCGCCCATGTGGAAACCGGCACGAAGGCGCGCCCTAATGCTCGTTTCTTCCAAGACGCGGACTGCATAGA

gD coding sequence

Ala Ala Arg Gly Tyr Thr Phe Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu Arg Leu Thr Tyr Leu

6x his:MgD coding seq

CACGCAGTACTACCCGAGGACACAAAGGCCATAGTCGACTACTGGTTTCATGCGGCCACGGGGCGTTCGTTCGCCCGT 880
GTGCGTCATGATGGGCGTCCCTCCGTGTGTTCCGGTATCAGCTGATGACCAAGTACGCGGTGCCCCCGCAGCAAGGCGGCA

gD coding sequence

Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala Ile Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro

6x his:MgD coding seq

ATTTTGAGGAGTCGAAGGCTACGAGCCGCCCTGCGCCCGATGGGGGTTCCCCCGGCCACCCGGCGACGACGAGGCC 960
TAAACTCCTCAGCTTCCCGATGCTCGGCGGACGCGGCTACCCCAAGGGGCGCGGTGGGCCCGCTGCTGCTCCGG

gD coding sequence

Tyr Phe Glu Glu Ser Lys Gly Tyr Glu Pro Pro Ala Ala Asp Gly Gly Ser Pro Ala Pro Pro Gly Asp Asp Glu Ala

6x his:MgD coding seq

FIG. 6D




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D






FIG. 8A




GAGCGGCGGCTGCGACATGCTGGCGCTGATCGCAGACCCGCAGGTGGGGCGCACGCTGTGGGAAGCGGTACGCCGGC
 400
 Ser Ala Ala Cys Asp Met Leu Ala Leu Ile Ala Asp Pro Gln Val Gly Arg Thr Leu Thr Glu Ala Val Arg Arg
 mature gD coding seq
 His-gD-LHRH fusion
 ACGCGCGCGGTACAACGCCACGGTCATATGTTACAAGATCGAGAGCGGTCGCCCGCGCTGTACTACATGGAGTAC
 480
 His Ala Arg Ala Tyr Asn Ala Thr Val Ile Thr Tyr lys Ile Glu Ser Gly Cys Ala Arg Pro Leu Tyr Tyr Met Glu Tyr
 His-gD-LHRH fusion
 ACCGAGTGCAGCCAGGAAGCACTTTGGGTACTGCCCGTACCGCACACCCCGTTTGGGACAGCTTCCTGCGCGGCTT
 560
 Thr Glu Cys Glu Pro Arg Lys His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Thr Asp Ser Phe Leu Ala Gly Phe
 His-gD-LHRH fusion
 CGCCTACCCACGGACGACGAGCTGGGACTGATTATGGCGCGCGCGCTCGTCTGAGGGCCAGTACCGACGCGCGC
 640
 Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile Met Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala
 His-gD-LHRH fusion
 TGTACATCGACGGCAGGTGCGCCTATACAGATTTCATGGTTTCGTCGCCGGGGACTGCTGGTTCTCGAAACTCGGC
 720
 Leu Tyr Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro Ala Gly Asp Cys Thr Phe Ser Lys Leu Gly
 His-gD-LHRH fusion

FIG. 8C



+

FIG. 8D



+




FIG. 9B




FIG. 9C




FIG. 9D




FIG. 10




FIG. 11A






FIG. 11B





Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	



+

700

■ Mature gd Sequence ———
The Gly Ala Cys His Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu Arg Leu Thr
gd-LHRH Coding Sequence

008

900

1000

— Mature gD Sequence —
 Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala Glu Gly Glu Pro Lys Pro Gly
 gD-LHRH Coding Sequence —

+



SEQUENCE LISTING

<110> Pfizer Products Inc

<120> FUSION PROTEINS COMPRISING CARRIERS THAT CAN INDUCE A
DUAL IMMUNE RESPONSE

<130> PC10202A

<140>

<141>

<150> N/A

<151> 1999-02-17

<160> 46

<170> PatentIn Ver. 2.1

<210> 1

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 1

catggaacac tggctcttatg gtctgcgtcc ggg

33

<210> 2

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 2

catggaacac tggctcttatg gtctgcgtcc ggg

33

<210> 3

<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 3
gatctggaac actggtctta tggctctgct cggggc

36

<210> 4
<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 4
gatcgcccg acgcagacca taagaccagt gttcca

36

<210> 5
<211> 76
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 5
gatccatgga gcactggtca tatggtctgc gtccgggtga acattggagc tacgggtctac
gccccgggtc catggc

60
76

<210> 6
<211> 76
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 6
tcgagccatg gacccggggc gtagaccgta gctccaatgt tcacccggac gcagaccata 60
tgaccagtgc tccatg 76

<210> 7
<211> 71
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 7
ggggaacact ggtcttatgg cttacggccg ggagagcatt ggagttacgg cctccgtcca 60
ggttccatgg c 71

<210> 8
<211> 75
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 8
tcgagccatg gaacctggac ggaggccgta actccaatgc tctcccggcc gtaagccata 60
agaccagtgt tcccc 75

<210> 9
<211> 71
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 9
gatccagagc actggtcata tggctgcgt ccgggtgaac attggagcta cggctctacgc 60
cccggggatc c 71

<210> 10
<211> 71
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 10
tcgaggatcc ccggggcgta gaccgtagct ccaatgttca ccgggacgca gaccatatga 60
ccagtgtct g 71

<210> 11
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 11
ggggaacact ggtcttatgg cttacggccg ggagagcatt ggagttacgg cctccgtcca 60
ggggatcc 68

<210> 12
<211> 72
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: SYNTHETIC
OLIGONUCLEOTIDE COMPRISING GNRH CODING SEQUENCE
AND CLONING ENDS

<400> 12
tcgaggatcc cctggacgga ggccgtaact ccaatgtctt ccgggccgta agccataaga 60

ccagtgttcc cc

72

<210> 13
<211> 10
<212> PRT
<213> GNRH AMINO ACID SEQUENCE

<400> 13
Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
1 5 10

<210> 14
<211> 328
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: part of
plasmid p9897-R

<400> 14
acgccagggt tttcccagtc acgacgttgt aaaacgacgg ccagtgagcg cgcgtaatac 60
gactcactat agggcgaatt ggagctccac cgcggtggcg gccgctctag aactagtgga 120
tccagagcac tggtcatatg gtctgcgtcc gggatgaacat tggagctacg gtctacgccc 180
cggggaacac tgggtcttatg gcttacggcc gggagagcat tggagttacg gcctccgtcc 240
aggttccatg ggctcgaggg ggggcccggg acccagcttt tgttcccttt agtgagggtt 300
aattgcgcgc ttggcgtaat atggtcat 328

<210> 15
<211> 40
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: GnRH tetramer

<400> 15
Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His Trp Ser Tyr Gly
1 5 10 15

Leu Arg Pro Gly Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His
20 25 30

Trp Ser Tyr Gly Leu Arg Pro Gly
35 40

5

EXPRESS MAIL NO. EL162820034US

<210> 16
<211> 1259
<212> DNA
<213> Bovine herpesvirus 1

<220>
<221> gene
<222> (1)..(1259)
<223> sequence encoding BHV-1 gD from clone
FlgD/Pots207(#79)

<400> 16
ccatggaggg gccgacattg gccgtgctgg gcgcgctgct cgcggttgcg gtaagcttgc 60
ctacacccgc gccgcgggtg acggtatacg tcgacccgcc ggcgtaaccg atgccgcgat 120
acaactaac tgaacgctgg cacactaccg ggcccatacc gtcgcccttc gcagacggcc 180
gogagcagcc cgtcgaggtg cgctacgcga cgagcgcggc ggcgtagcac atgctggcgc 240
tgatcgaga cccgcaggtg gggcgcacgc tgtgggaagc ggtacgccgg cacgcgcgcg 300
cgtacaacgc caccgtcata tggtagaaga tcgagagcgg gtgcgccggg ccgctgtact 360
acatggagta caccgagtgc gagcccagga agcactttgg gtactgccgc taccgcacac 420
ccccgttttg ggacagcttc ctggcgggct tcgcctaccc cacggacgac gagctgggac 480
tgattatggc ggcgcccgcg cggctcgtcg agggccagta ccgacgcgcg ctgtacatcg 540
acggcacggg cgccataaca gatttcatgg tttcgtgccc ggccggggac tgctggttct 600
cgaaactcgg cgcggctcgc gggtagacct ttggcgcgtg cttcccggcc cgggattacg 660
agcaaaagaa ggttctgcgc ctgacgtatc tcacgcagta ctaccgcgag gaggcacaca 720
aggccatagt cgactactgg ttcatgcgcc acgggggcgt cgttcgcgcg tattttgagg 780
agtcgaaggg ctacgagccg ccgcctgcgc ccgatggggg ttcccccgcg ccaccggcg 840
acgacgaggc ccgcgaggat gaaggggaga ccgaggacgg ggcagccggg cgggagggca 900
acggcgggcc cccaggaccc gaaggcgacg gcgagagtca gacccccgaa gccaacggag 960
gcgccgaggg cgagccgaaa cccggccccca gccccgacgc cgaccgcccc gaaggctggc 1020
cgagcctcga agccatcacg cccccccgc ccgccccgc tacgcccgt cgagctccgg 1080
acgtgtttc ggtttctggt ggtatcggtg tcgctgctgc tgctatcgt tcggttgctg 1140
ctgctgctgc tgggtgcttac ttcgtttata ttgctgctgc tgggtgctgg ccgctgccgc 1200
gtaaaccgaa aaaactgccg gctttcggtg acgttaacta cagtgtctct cggggttga 1259

<210> 17
<211> 418
<212> PRT
<213> Bovine herpesvirus 1

<220>
<221> PEPTIDE
<222> (1)..(418)
<223> BHV-1gD encoded by clone FlgD/Pots207nco(#79)

<400> 17

Met Glu Gly Pro Thr Leu Ala Val Leu Gly Ala Leu Leu Ala Val Ala
 1 5 10 15
 Val Ser Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro
 20 25 30
 Pro Ala Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr
 35 40 45
 Thr Gly Pro Ile Pro Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val
 50 55 60
 Glu Val Arg Tyr Ala Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu
 65 70 75 80
 Ile Ala Asp Pro Gln Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg
 85 90 95
 His Ala Arg Ala Tyr Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser
 100 105 110
 Gly Cys Ala Arg Pro Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro
 115 120 125
 Arg Lys His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp
 130 135 140
 Ser Phe Leu Ala Gly Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu
 145 150 155 160
 Ile Met Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala
 165 170 175
 Leu Tyr Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu
 180 185 190
 Pro Ala Gly Asp Cys Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr
 195 200 205
 Thr Phe Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val
 210 215 220
 Leu Arg Leu Thr Tyr Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys
 225 230 235 240
 Ala Ile Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro
 245 250 255

Tyr Phe Glu Glu Ser Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly
 260 265 270
 Gly Ser Pro Ala Pro Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly
 275 280 285
 Glu Thr Glu Asp Gly Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro
 290 295 300
 Gly Pro Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly
 305 310 315 320
 Ala Glu Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro
 325 330 335
 Glu Gly Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro
 340 345 350
 Ala Thr Pro Ala Arg Ala Pro Asp Ala Val Ser Val Ser Val Gly Ile
 355 360 365
 Gly Ile Ala Ala Ala Ala Ile Ala Cys Val Ala Ala Ala Ala Ala Gly
 370 375 380
 Ala Tyr Phe Val Tyr Ile Arg Arg Arg Gly Ala Gly Pro Leu Pro Arg
 385 390 395 400
 Lys Pro Lys Lys Leu Pro Ala Phe Gly Asn Val Asn Tyr Ser Ala Leu
 405 410 415
 Pro Gly

<210> 18

<211> 1405

<212> DNA

<213> Bovine herpesvirus 1

<220>

<221> gene

<222> (1)..(1405)

<223> BHV-1 gD from GenBank Accession No. M59846.

<400> 18

gggccgcagc cccggctggg tatatatccc cgacgggcga ctagagatac actcgccccg 60
 cgcggctgct gcgagcgggc gaacatgcaa gggccgacat tggccgtgct gggcgcgctg 120
 ctgcgcgttg cggtagctt gcctacacc gcgccgcggg tgacggtata cgtcgacccg 180

3403030

ccggcggtacc cgatgccgcg atacaactac actgaacgct ggcacactac cgggcccata 240
ccgtcgccct tcgcagacgg ccgcgagcag cccgtcgagg tgcgctacgc gacgagcgcg 300
gcggcggtgcg acatgctggc gctgatcgca gaccgcgagg tggggcgcac gctgtgggaa 360
gcgggtacgcc ggcacgcgcg ccgcgtacaac gccacgggtca tatggtacaa gatcgagagc 420
gggtgcgccc ggccgctgta ctacatggag tacaccgagt gcgagcccag gaagcacttt 480
gggtactgcc gctaccgcac acccccgttt tgggacagct tcctggcggg cttcgcctac 540
cccacggacg acgagctggg actgattatg gcggcgcccg cgcggctcgt cgagggccag 600
taccgacgcg cgctgtacat cgacggcacg gtcgcctata cagatttcat ggtttcgctg 660
ccggccggggg actgctggtt ctcgaaactc ggcgcggtc gcgggtacac ctttggcgcg 720
tgcttcccgg ccggggatta cgagcaaaaag aaggttctgc gcctgacgta tctcacgcag 780
tactaccgcg aggaggcaca caaggccata gtcgactact ggttcatgcg ccacgggggc 840
gtcgttccgc cgtattttga ggagtogaag ggctacgagc cgccgcctgc cgccgatggg 900
ggttcccccg cgccaccgag cgacgacgag gcccgcgagg atgaagggga gaccgaggac 960
ggggcagccg ggccgggagg caacggcggc cccccaggac ccgaaggcga cggcgagagt 1020
cagacccccg aagccaacgg aggcgcgag ggcgagccga aaccggccc cagccccgac 1080
gccgaccgcc ccgaaggctg gccgagcctc gaagccatca cgcaccccc gcccgcccc 1140
gctacgcccg cgcccccgga cgccgtgccg gtcagcgctc ggatcggcac tgcggctgcg 1200
gcgatcgctg cgttgccgc cgccgcgcgc ggcgcgtaact tcgtctatac gcgccggcg 1260
ggtgcgggtc cgctgcccag aaagccaaaa aagctgcgg cctttggcaa cgtcaactac 1320
agcgcgctgc ccgggtgagc ggcctaggcc ctcccccgac cgcccccttt gctcctagcc 1380
ccggctcctg ccgagccgcg cgggg 1405

<210> 19

<211> 417

<212> PRT

<213> Bovine herpesvirus 1

<220>

<221> PEPTIDE

<222> (1)..(417)

<223> BHV-1 gD encoded by GenBank Accession No. M59846.

<400> 19

Met Gln Gly Pro Thr Leu Ala Val Leu Gly Ala Leu Leu Ala Val Ala
1 5 10 15

Val Ser Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro
20 25 30

Pro Ala Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr
35 40 45

Thr Gly Pro Ile Pro Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val
50 55 60

Glu Val Arg Tyr Ala Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu
65 70 75 80

Ile Ala Asp Pro Gln Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg
85 90 95

His Ala Arg Ala Tyr Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser
100 105 110

Gly Cys Ala Arg Pro Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro
115 120 125

Arg Lys His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp
130 135 140

Ser Phe Leu Ala Gly Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu
145 150 155 160

Ile Met Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala
165 170 175

Leu Tyr Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu
180 185 190

Pro Ala Gly Asp Cys Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr
195 200 205

Thr Phe Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val
210 215 220

Leu Arg Leu Thr Tyr Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys
225 230 235 240

Ala Ile Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro
245 250 255

Tyr Phe Glu Glu Ser Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly
260 265 270

Gly Ser Pro Ala Pro Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly
275 280 285

Glu Thr Glu Asp Gly Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro
290 295 300

Gly Pro Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly
305 310 315 320

Ala Glu Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro
325 330 335

Glu Gly Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro
340 345 350

Ala Thr Pro Ala Ala Pro Asp Ala Val Pro Val Ser Val Gly Ile Gly
355 360 365

Ile Ala Ala Ala Ala Ile Ala Cys Val Ala Ala Ala Ala Gly Ala
370 375 380

Tyr Phe Val Tyr Thr Arg Arg Arg Gly Ala Gly Pro Leu Pro Arg Lys
385 390 395 400

Pro Lys Lys Leu Pro Ala Phe Gly Asn Val Asn Tyr Ser Ala Leu Pro
405 410 415

Gly

<210> 20
<211> 1218
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence from
pQE-tmgD encoding a tmgD.

<400> 20

```
ctcgagaaat cataaaaaat ttatttgctt tgtgagcgga taacaattat aatagattca 60
attgtgagcg gataacaatt tcacacagaa ttcattaaag aggagaaatt aactatgaga 120
ggatctcacc atcaccatca ccatacggat ccgcatgcca tgagcttgcc tacaccgcg 180
ccgcgggtga cggatatacgt cgaccgcgcg gcgatcccgga tgccgcgata caactacact 240
gaacgctggc aactaccgg gccataaccg tcgcccttcg cagacggccg cgagcagccc 300
gtcgaggtgc gctacgcgac gagcgcgcg gcggtgcgaca tgctggcgct gatcgagac 360
ccgcaggtgg ggcgcacgct gtgggaagcg gtacgccggc acgcgcgcgc gtacaacgcc 420
acggtcatat ggtacaagat cgagagcggg tgcgccggc cgctgtacta catggagtac 480
accgagtgcg agcccaggaa gcactttggg tactgccgct accgcacacc ccggttttg 540
gacagcttcc tggcgggctt cgctacccc acggacgacg agctgggact gattatggcg 600
gcgcccgcgc ggctcgtcga gggccagtac cgacgcgcgc tgtacatoga cggcacggtc 660
gcctatacag atttcatggt ttcgctgccg gccggggact gctggttctc gaaactcggc 720
gcggctcgcg ggtacacctt tggcgcgtgc ttcccgcccc gggattacga gcaaaagaag 780
gttctgcgcc tgacgtatct cacgcagtac taccgcagg aggcacacaa ggccatagtc 840
gactactggt tcatgcgcca cgggggcgct gttccgccgt attttgagga gtcgaagggc 900
tacgagccgc cgctgcccgc cgatgggggt tccccgcgc caccggcgca cgacgaggcc 960
cgcgaggatg aaggggagac cgaggacggg gcagccgggc gggagggcaa cggcggcccc 1020
ccaggacccg aaggcgacgg cgagagtcag acccccgaag ccaacggagg cgccgagggc 1080
```

gagccgaaac cccgccccag ccccgacgcc gaccgccccg aaggctggcc gagcctcgaa 1140
gccatcagc accccccgcc cccccccgct acgcccgtc gagctcggta ccccgggtcg 1200
acctgcagcc aagcttaa 1218

<210> 21
<211> 367
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: tmgD encoded
by pQE-tmgD.

<400> 21
Met Arg Gly Ser His His His His His Thr Asp Pro His Ala Met
1 5 10 15
Ser Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro
20 25 30
Ala Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr
35 40 45
Gly Pro Ile Pro Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val Glu
50 55 60
Val Arg Tyr Ala Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu Ile
65 70 75 80
Ala Asp Pro Gln Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg His
85 90 95
Ala Arg Ala Tyr Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser Gly
100 105 110
Cys Ala Arg Pro Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro Arg
115 120 125
Lys His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp Ser
130 135 140
Phe Leu Ala Gly Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile
145 150 155 160
Met Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala Leu
165 170 175

Tyr Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro
 180 185 190
 Ala Gly Asp Cys Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr Thr
 195 200 205
 Phe Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu
 210 215 220
 Arg Leu Thr Tyr Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala
 225 230 235 240
 Ile Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro Tyr
 245 250 255
 Phe Glu Glu Ser Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly Gly
 260 265 270
 Ser Pro Ala Pro Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly Glu
 275 280 285
 Thr Glu Asp Gly Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro Gly
 290 295 300
 Pro Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala
 305 310 315 320
 Glu Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu
 325 330 335
 Gly Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala
 340 345 350
 Thr Pro Ala Arg Ala Arg Tyr Pro Gly Ser Thr Cys Ser Gln Ala
 355 360 365

<210> 22

<211> 1360

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: portion of
 pQE-GnRH:gD, including sequence encoding
 4GnRH-tmgD.

<400> 22

```

ctcgagaaat cataaaaaat ttatttgctt tgtgagcgga taacaattat aatagattca 60
attgtgagcg gataacaatt tcacacagaa ttcattaaag aggagaaatt aactatgaga 120
ggatctcacc atcaccatca ccatacggat ccgcatgcca tggatccaga gcaactggtca 180
tatggtctgc gtcggggtga acattggagc tacggtctac gccccgggga acactggtct 240
tatggcttac ggccgggaga gcattggagt tacggcctcc gtccagggtc catgagcttg 300
cctacacccg cgccgcggtg gacggtatac gtcgacccgc cggcgtaccc gatgccgcga 360
tacaactaca ctgaacgctg gcacactacc gggcccatac cgtcgccctt cgcagacggc 420
cgcgagcagc ccgtcgaggt gcgctacgag acgagcgcgg cggcgtgcga catgctggcg 480
ctgatcgagc acccgcaggt ggggcgacg ctgtgggaag cggtagccg gcacgcgcgc 540
gcgtacaacg ccacggtcat atggtacaag atcgagagcg ggtgcgccc gccgctgtac 600
tacatggagt acaccgagtg cgagcccagg aagcactttg ggtactgcc ctaccgcaca 660
cccccgtttt gggacagctt cctggcgggc ttgcctacc ccacggacga cgagctggga 720
ctgattatgg cggcgccgcg gcggctcgtc gagggccagt accgacgcgc gctgtacatc 780
gacggcacgg tcgcctatac agatttcatg gtttcgctgc cggccgggga ctgctggttc 840
tcgaaactcg gcgcggctcg cgggtacacc tttggcgctg gcttcccgc cggggattac 900
gagcaaaaga aggttctgcg cctgacgtat ctcacgcagt actaccgca ggaggcacac 960
aaggccatag tcgactactg gttcatgcgc cacgggggag tcgttccgc gtattttgag 1020
gagtcgaagg gctacgagcc gcgcctgcc gccgatggg gttccccgc gccaccggc 1080
gacgacgagg ccgcgagga tgaaggggag accgaggagc gggcagccg gcgggagggc 1140
aacggcgcc ccccaggacc cgaaggcgac ggcgagagtc agacccccga agccaacgga 1200
ggcgccgagg gcgagccgaa acccgcccc agccccgag ccgaccgcc cgaaggctgg 1260
ccgagcctcg aagccatcac gcacccccg cccgccccg ctacgcccgc tcgagctcgg 1320
taccgccggt cgacctgag ccaagcttaa ttagctgagc 1360

```

<210> 23

<211> 411

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 4GnRH-tmgD
encoded by pQE-GnRH:gD.

<400> 23

```

Met Arg Gly Ser His His His His His Thr Asp Pro His Ala Met
  1             5             10             15

```

```

Asp Pro Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His Trp Ser
      20             25             30

```

```

Tyr Gly Leu Arg Pro Gly Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
      35             40             45

```

```

Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Ser Met Ser Leu Pro Thr
      50             55             60

```

```

Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro Ala Tyr Pro Met

```


65		70		75		80
Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr Gly Pro Ile Pro						
	85		90		95	
Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val Glu Val Arg Tyr Ala						
	100		105		110	
Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu Ile Ala Asp Pro Gln						
	115		120		125	
Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg His Ala Arg Ala Tyr						
	130		135		140	
Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser Gly Cys Ala Arg Pro						
	145		150		155	160
Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro Arg Lys His Phe Gly						
		165		170		175
Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp Ser Phe Leu Ala Gly						
	180		185		190	
Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile Met Ala Ala Pro						
	195		200		205	
Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala Leu Tyr Ile Asp Gly						
	210		215		220	
Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro Ala Gly Asp Cys						
	225		230		235	240
Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr Thr Phe Gly Ala Cys						
		245		250		255
Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu Arg Leu Thr Tyr						
	260		265		270	
Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala Ile Val Asp Tyr						
	275		280		285	
Trp Phe Met Arg His Gly Gly Val Val Pro Pro Tyr Phe Glu Glu Ser						
	290		295		300	
Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly Gly Ser Pro Ala Pro						
	305		310		315	320
Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly Glu Thr Glu Asp Gly						

325

330

335

Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro Gly Pro Glu Gly Asp
 340 345 350

Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala Glu Gly Glu Pro
 355 360 365

Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu Gly Trp Pro Ser
 370 375 380

Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala Thr Pro Ala Arg
 385 390 395 400

Ala Arg Tyr Pro Gly Ser Thr Cys Ser Gln Ala
 405 410

<210> 24

<211> 1360

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: portion of
 pQE-gD:GnRH, including sequence coding tmgD-4GnRH.

<400> 24

ctcgagaaat cataaaaaat ttatttgctt tgtgagcgga taacaattat aatagattca 60
 attgtgagcg gataacaatt tcacacagaa ttcattaaag aggagaaatt aactatgaga 120
 ggatctcacc atcaccatca ccatacggat ccgcatgcca tgagcttgcc tacaccgcg 180
 ccgcggtga cggatatacgt cgaccgcgcg gcgtacccga tgccgcgata caactacact 240
 gaacgctggc aactaccgg gccataccg tcgcccttcg cagacggcgc cgagcagccc 300
 gtcgaggtgc gctacgcgac gagcgcggcg gcgtgcgaca tgctggcgct gatcgcagac 360
 ccgcaggtgg ggcgcacgct gtgggaagcg gtacgcgcgc acgcgcgcgc gtacaacgcc 420
 acggtcatat ggtacaagat cgagagcggg tcgcgccggc cgctgtacta catggagtac 480
 accgagtgcg agcccaggaa gcactttggg tactgccgct accgcacacc ccggttttgg 540
 gacagcttcc tggcgggctt cgctacccc acggaacgac agctgggact gattatggcg 600
 gcgcccgcgc ggctcgtcga gggccagtac cgacgcgcgc tgtacatcga cggcacggtc 660
 gcctatacag atttcatggt ttcgctgcg gccggggact gctggttctc gaaactcggc 720
 gcggctcgcg ggtacacctt tggcgcgctc ttcccggccc gggattacga gcaaaagaag 780
 gttctgcgcc tgacgtatct cacgcagtac taccgcagc aggcacacaa ggccatagtc 840
 gactactggt tcatgcgcca cggggcgctc gttccgcgct attttgagga gtcgaaggcg 900
 tacgagccgc cgctgcgcgc cgatgggggt tccccgcgc caccgcgcga cgacgaggcc 960
 cgcgaggatg aaggggagac cgaggacggg gcagccgggc gggagggcaa cggcggcccc 1020
 ccaggacccg aaggcgacgg cgagagtcag acccccgaag ccaacggagg cgccgagggc 1080
 gagccgaaac ccggccccag ccccgacgcc gaccgcccc aaggctggcc gagcctcgaa 1140
 gccatcacgc accccccgcg cgccccgct acgcccgcgc gagctccaga gcactggtca 1200

tatggtctgc gtcgggtga acattggagc tacggtctac gccccgggga acactggtct 1260
 tatggcttac ggccgggaga gcattggagt tacggcctcc gtccaggttg aagcttaatt 1320
 agctgagctt ggactcctgt tgatagatcc agtaatgacc 1360

<210> 25

<211> 398

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: tmgD-4GmRH
 encoded by pQE-gD:GmRH.

<400> 25

Met Arg Gly Ser His His His His His Thr Asp Pro His Ala Met
 1 5 10 15

Ser Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro
 20 25 30

Ala Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr
 35 40 45

Gly Pro Ile Pro Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val Glu
 50 55 60

Val Arg Tyr Ala Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu Ile
 65 70 75 80

Ala Asp Pro Gln Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg His
 85 90 95

Ala Arg Ala Tyr Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser Gly
 100 105 110

Cys Ala Arg Pro Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro Arg
 115 120 125

Lys His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp Ser
 130 135 140

Phe Leu Ala Gly Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile
 145 150 155 160

Met Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala Leu
 165 170 175

Tyr Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro
180 185 190

Ala Gly Asp Cys Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr Thr
195 200 205

Phe Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu
210 215 220

Arg Leu Thr Tyr Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala
225 230 235 240

Ile Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro Tyr
245 250 255

Phe Glu Glu Ser Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly Gly
260 265 270

Ser Pro Ala Pro Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly Glu
275 280 285

Thr Glu Asp Gly Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro Gly
290 295 300

Pro Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala
305 310 315 320

Glu Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu
325 330 335

Gly Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala
340 345 350

Thr Pro Ala Arg Ala Pro Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
355 360 365

Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His Trp Ser Tyr Gly
370 375 380

Leu Arg Pro Gly Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
385 390 395

<210> 26

<211> 1441

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: portion of
pQE-GnRH:gD:GnRH, including encoding
4GnRH-tmgD-4GnRH

<400> 26

ctcgagaaat cataaaaaat ttatttgctt tgtgagcggg taacaattat aatagattca 60
attgtgagcg gataacaatt tcacacagaa ttcattaaag aggagaaatt aactatgaga 120
ggatctcacc atcaccatca ccatacggat ccgcatgccg tggatccaga gacttggtca 180
tatggtctgc gtccgggtga acattggagc tacggtctac gcccggggga aacttggtct 240
tatggcttac ggccgggaga gcattggagt tacggcctcc gtccaggttc catgagcttg 300
cctacaccgg cgccgggggt gacggtatcc gtcgaccggc cggcgtagcc gatgccgcga 360
tacaactaca ctgaacgctg gcacactacc gggcccatcc cgtcgccctt cgcagacggc 420
cgcgagcagc ccgtcgaggt gcgctacggc acgagcgcgg cggcgtagcc catgctggcg 480
ctgatcgagc acccgaggt gggcgagcag ctgtgggaag cggtagcccg gcacgcgcgc 540
gcgtacaacg ccacggtcat atggtacaag atcgagagcg ggtgcgcccg gccgctgtac 600
tacatggagt acaccgagtg cgagcccagg aagcactttg ggtactgccg ctaccgcaca 660
ccccggtttt gggacagctt cctggcgggc ttgcgctacc ccacggacga cgagctggga 720
ctgattatgg cggcgcccg cgggctcgtc gagggccagt accgacgcgc gctgtacatc 780
gacggcacgg tcgctatac agatttcatg gtttcgctgc cggccgggga ctgctgggtc 840
tcgaaactcg gcgcggtcg cgggtacacc tttggcgctg gtttcccgcc cggggattac 900
gagcaaaaga aggttctgcg cctgacgtat ctcacgcagt actaccgca ggaggcacac 960
aaggccatag tcgactactg gttcatgcgc cacggggggc tcgttccgcc gtattttgag 1020
gagtcgaagg gctacgagcc gccgcctgcc gccgatgggg gttccccgc gccaccggc 1080
gacgacgagg cccgcgagga tgaaggggag accgaggacg gggcagccgg gcgggagggc 1140
aacggcggcc cccaggacc cgaaggcgac ggcgagagtc agacccccga agccaacgga 1200
ggcgccgagg gcgagccgaa acccgggccc agccccgac cgcaccgcc cgaaggctgg 1260
ccgagcctcg aagccatcac gcaccccccg cccgcccccg ctacgcccgc tcgagctcca 1320
gagcactggc catatggtct gcgtccgggt gaacattgga gctaoggtct acgccccggg 1380
gaacactggc cttatggctt acggccggga gagcattgga gttacggcct ccgtccaggt 1440
t 1441

<210> 27

<211> 442

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
4GnRH-tmgD-4GnRH encoded by pQE-GnRH:gD:GnRH

<400> 27

Met Arg Gly Ser His His His His His His Thr Asp Pro His Ala Met
1 5 10 15
Asp Pro Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His Trp Ser
20 25 30

Tyr Gly Leu Arg Pro Gly Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
35 40 45

Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Ser Met Ser Leu Pro Thr
50 55 60

Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro Ala Tyr Pro Met
65 70 75 80

Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr Gly Pro Ile Pro
85 90 95

Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val Glu Val Arg Tyr Ala
100 105 110

Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu Ile Ala Asp Pro Gln
115 120 125

Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg His Ala Arg Ala Tyr
130 135 140

Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser Gly Cys Ala Arg Pro
145 150 155 160

Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro Arg Lys His Phe Gly
165 170 175

Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp Ser Phe Leu Ala Gly
180 185 190

Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile Met Ala Ala Pro
195 200 205

Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala Leu Tyr Ile Asp Gly
210 215 220

Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro Ala Gly Asp Cys
225 230 235 240

Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr Thr Phe Gly Ala Cys
245 250 255

Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu Arg Leu Thr Tyr
260 265 270

Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala Ile Val Asp Tyr
275 280 285

Trp Phe Met Arg His Gly Gly Val Val Pro Pro Tyr Phe Glu Glu Ser
 290 295 300
 Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly Gly Ser Pro Ala Pro
 305 310 315 320
 Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly Glu Thr Glu Asp Gly
 325 330 335
 Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro Gly Pro Glu Gly Asp
 340 345 350
 Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala Glu Gly Glu Pro
 355 360 365
 Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu Gly Trp Pro Ser
 370 375 380
 Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala Thr Pro Ala Arg
 385 390 395 400
 Ala Pro Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His Trp Ser
 405 410 415
 Tyr Gly Leu Arg Pro Gly Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
 420 425 430
 Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
 435 440

<210> 28
 <211> 1079
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: portion of
 pCMV-tgD, including sequence encoding a truncated
 gD

<400> 28
 gatatcatgc aggggcccgc attggccgtg ctgggcgcgc tgctcgccgt tgcggttaagc 60
 ttgcctacac ccgcgcgcgc ggtgacggtg tacgtcgacc cgccggcgta cccgatgccg 120
 cgatacaact aactgaacg ctggcacact accgggcccga taccgtcgcc cttcgagac 180
 ggccgcgagc agcccgctga ggtgcgtac gcgacgagc cgccggcggtg cgacatgctg 240
 gcgctgatcg cagaccgcga ggtggggcgc acgctgtggg aagcggtacg ccggcacgcg 300

cgcgcggtaca acgccacggt catatggtac aagatcgaga gcgggtgcgc ccggccgctg 360
 tactacatgg agtacaccga gtgcgagccc aggaagcact ttgggtactg ccgctaccgc 420
 acacccccgt tttgggacag ctctctggcg ggcttcgcct accccacgga cgacgagctg 480
 ggactgatta tggcgggcgcc cgcgcggtc gtcgagggcc agtaccgacg cgcgctgtac 540
 atcgacggca cggtcgccta tacagatttc atggtttcgc tgccggccgg ggactgctgg 600
 ttctcgaaac tcggcgcggc tcgcggttac acctttggcg cgtgcttccc ggcccgggat 660
 tacgagcaaa agaaggttct gcgcctgacg tatctcacgc agtactaccc gcaggaggca 720
 cacaaggcca tagtcgacta ctggttcatg cgccacgggg gcgtcgttcc gccgtatatt 780
 gaggagtcga agggctacga gccgcgcct gccgccgatg ggggttcccc cgcgccaccc 840
 ggcgacgacg agggccgcga ggatgaagg gagaccgagg acggggcagc cgggcgggag 900
 ggcaacggcg gccccccagg acccgaaggc gacggcgaga gtcagacccc cgaagccaac 960
 ggaggcgccg agggcgagcc gaaacccggc ccagccccg acgccgaccg ccccgaggct 1020
 ggccgagcct cgaagccatc acgcaccccc cgcccgcccc cgctacgcc tgaggtacc 1079

<210> 29

<211> 353

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: truncated gD
encoded by pCMV-tgD

<400> 29

Met Gln Gly Pro Thr Leu Ala Val Leu Gly Ala Leu Leu Ala Val Ser
1 5 10 15

Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro Ala
20 25 30

Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr Gly
35 40 45

Pro Ile Pro Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val Glu Val
50 55 60

Arg Tyr Ala Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu Ile Ala
65 70 75 80

Asp Pro Gln Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg His Ala
85 90 95

Arg Ala Tyr Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser Gly Cys
100 105 110

Ala Arg Pro Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro Arg Lys
115 120 125

22

EXPRESS MAIL NO. EL16282003405

His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp Ser Phe
130 135 140

Leu Ala Gly Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile Met
145 150 155 160

Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala Leu Tyr
165 170 175

Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro Ala
180 185 190

Gly Asp Cys Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr Thr Phe
195 200 205

Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu Arg
210 215 220

Leu Thr Tyr Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala Ile
225 230 235 240

Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro Tyr Phe
245 250 255

Glu Glu Ser Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly Gly Ser
260 265 270

Pro Ala Pro Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly Glu Thr
275 280 285

Glu Asp Gly Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro Gly Pro
290 295 300

Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala Glu
305 310 315 320

Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu Gly
325 330 335

Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala Thr
340 345 350

Pro

<210> 30

<211> 1241
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: portion of
 pCMV-gD:GnRH, including sequence encoding a
 tgD-4GnRH fusion protein

<400> 30

```

gcggccgcaa gatatcatgc aggggcccgc attggccgtg ctggggcgcg tgctcgccgt 60
tgccgtaagc ttgcctacac ccgcgcgcgc ggtgacggtg tacgtcgacc cgccggcgta 120
cccgatgccg cgatacaact acactgaacg ctggcacact accgggcccc taccgtcgcc 180
cttcgcagac ggccgcgagc agcccgtcga ggtgcgctac gcgacgagcg cggcggcggtg 240
cgacatgctg gcgctgatcg cagaccgcga ggtggggcgc acgctgtggg aagcggtagc 300
ccggcacgcg cgcgcgtaca acgccacggt catatggtac aagatcgaga gcgggtgcgc 360
ccggccgctg tactacatgg agtacaccga gtgcgagccc aggaagcact ttgggtactg 420
ccgctaccgc acacccccgt tttgggacag cttcctggcg ggcttcgcct accccacgga 480
cgacgagctg ggactgatta tggcggcgcc cgcgcggctc gtcgagggcc agtaccgacg 540
cgcgctgtac atcgacggca cggtcgccta tacagatttc atggtttcgc tgccggcccg 600
ggactgctgg ttctcgaaac tcggcgcggc tcggcggtac acctttggcg cgtgcttccc 660
ggcccgggat tacgagcaaa agaaggttct gcgcctgacg tatctcacgc agtactaccc 720
gcaggaggca cacaaggcca tagtcgacta ctggttcatg cggcacgggg gcgtcgttcc 780
gccgtatttt gaggagtcga agggctacga gccgcgcct gccgccgatg ggggttcccc 840
cgcgccaccc ggcgacgacg agggccgcga ggatgaaggg gagaccgagg acggggcagc 900
cgggcgggag ggcaacggcg gccccccagg acccgaaggc gacggcgaga gtcagacccc 960
cgaagccaac ggaggcgcgc agggcgagcc gaaacccggc ccagccccc acgcccagcg 1020
ccccgaaggc tggccgagcc tcgaagccat caccgacccc ccgcccggcc ccgctacgcc 1080
cgctcgagct ccagagcact ggtcatatgg tctgcgtccg ggtgaacatt ggagctacgg 1140
tctacgcccc ggggaacact ggtcttatgg cttacggccg ggagagcatt ggagttacgg 1200
cctccgtcca ggttgaagct gggatactag tgagcggccg c 1241

```

<210> 31
 <211> 397
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: tgD-4GnRH
 fusion protein encoded by pCMV-gD:GnRH

<400> 31

```

Met Gln Gly Pro Thr Leu Ala Val Leu Gly Ala Leu Leu Ala Val Ser
  1              5              10              15

Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro Ala
          20              25              30

```

Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr Gly
35 40 45

Pro Ile Pro Ser Pro Phe Ala Asp Gly Arg Glu Gln Pro Val Glu Val
50 55 60

Arg Tyr Ala Thr Ser Ala Ala Ala Cys Asp Met Leu Ala Leu Ile Ala
65 70 75 80

Asp Pro Gln Val Gly Arg Thr Leu Trp Glu Ala Val Arg Arg His Ala
85 90 95

Arg Ala Tyr Asn Ala Thr Val Ile Trp Tyr Lys Ile Glu Ser Gly Cys
100 105 110

Ala Arg Pro Leu Tyr Tyr Met Glu Tyr Thr Glu Cys Glu Pro Arg Lys
115 120 125

His Phe Gly Tyr Cys Arg Tyr Arg Thr Pro Pro Phe Trp Asp Ser Phe
130 135 140

Leu Ala Gly Phe Ala Tyr Pro Thr Asp Asp Glu Leu Gly Leu Ile Met
145 150 155 160

Ala Ala Pro Ala Arg Leu Val Glu Gly Gln Tyr Arg Arg Ala Leu Tyr
165 170 175

Ile Asp Gly Thr Val Ala Tyr Thr Asp Phe Met Val Ser Leu Pro Ala
180 185 190

Gly Asp Cys Trp Phe Ser Lys Leu Gly Ala Ala Arg Gly Tyr Thr Phe
195 200 205

Gly Ala Cys Phe Pro Ala Arg Asp Tyr Glu Gln Lys Lys Val Leu Arg
210 215 220

Leu Thr Tyr Leu Thr Gln Tyr Tyr Pro Gln Glu Ala His Lys Ala Ile
225 230 235 240

Val Asp Tyr Trp Phe Met Arg His Gly Gly Val Val Pro Pro Tyr Phe
245 250 255

Glu Glu Ser Lys Gly Tyr Glu Pro Pro Pro Ala Ala Asp Gly Gly Ser
260 265 270

Pro Ala Pro Pro Gly Asp Asp Glu Ala Arg Glu Asp Glu Gly Glu Thr
275 280 285

Glu Asp Gly Ala Ala Gly Arg Glu Gly Asn Gly Gly Pro Pro Gly Pro
 290 295 300

Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala Glu
 305 310 315 320

Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu Gly
 325 330 335

Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala Thr
 340 345 350

Pro Ala Arg Ala Pro Glu His Trp Ser Tyr Gly Leu Arg Pro Gly Glu
 355 360 365

His Trp Ser Tyr Gly Leu Arg Pro Gly Glu His Trp Ser Tyr Gly Leu
 370 375 380

Arg Pro Gly Glu His Trp Ser Tyr Gly Leu Arg Pro Gly
 385 390 395

<210> 32

<211> 120

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sequence
 encoding a GnRH tetramer

<400> 32

gagcactggt catatggtct gcgtccgggt gaacattgga gctacggtct acgccccggg 60
 gaacactggt cttatggcctt acggccggga gagcattgga gttacggcct ccgtccaggt 120

<210> 33

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sequence
 encoding a GnRH monomer

<400> 33

gagcactggt catatggtct gcgtccgggt

30

<210> 34
 <211> 1179
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sequence
 encoding a 4GnRH-tmgD fusion protein

<400> 34

```

gagcactggt catatggtct gcgtccgggt gaacattgga gctacggtct acgccccggg 60
gaacactggt cttatggctt acggccggga gagcattgga gttacggcct ccgtccaggt 120
tccatgagct tgctacacc cgcgcgcggg gtgacggtat acgtcgaccc gccggcgtag 180
ccgatgccgc gatacaacta cactgaacgc tggcacacta ccggggccat accgtcgccc 240
ttcgcagacg gccgcgagca gcccgtagag gtgcgctacg cgacgagcgc ggcggcgtgc 300
gacatgctgg cgtgatcgc agaccgcgag gtggggcgca cgctgtggga agcggtagcg 360
cggcacgcgc gcgcgtacaa cgccacgggtc atatggtaca agatcgagag cgggtgcgcc 420
cggccgctgt actacatgga gtacaccgag tgcgagccca ggaagcactt tgggtactgc 480
cgctaccgca caccgccgtt ttgggacagc ttcttgggcg gcttcgccta cccacaggac 540
gacgagctgg gactgattat ggcggcgccc gcgcggctcg tcgagggcca gtaccgacgc 600
gcgctgtaca tcgacggcac ggctgcctat acagatttca tggtttcgct gccggccggg 660
gactgctggt tctcgaact cggcgcggtt cgcgggtaca cctttggcgc gtgcttccc 720
gcccgggatt acgagcaaaa gaagggtctg cgctgacgt atctcacgca gtactaccg 780
caggaggcac acaaggccat agtcgactac tggttcatgc gccacggggg cgtcgttccg 840
ccgtattttg aggagtcgaa gggctacgag ccgccgcctg ccgcgcgatg gggttcccc 900
gcgccacccg gcgacgacga gggccgcgag gatgaagggg agaccgagga cggggcagcc 960
gggcgggagg gcaacggcgg cccccagga ccgaaggcg acggcgagag tcagaccccc 1020
gaagccaacg gaggcgccga gggcgagccg aaacccggcc ccagcccoga cgcgcagcgc 1080
cccgaaggct ggccgagcct cgaagccatc acgcaccccc cgcgcgcccc cgctacgccc 1140
gctcgagctc ggtaccccg gtcgacctgc agccaagct 1179
  
```

<210> 35
 <211> 340
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: a truncated
 mature BHV-1 gD

<400> 35

```

Leu Pro Thr Pro Ala Pro Arg Val Thr Val Tyr Val Asp Pro Pro Ala
1           5           10          15
  
```

```

Tyr Pro Met Pro Arg Tyr Asn Tyr Thr Glu Arg Trp His Thr Thr Gly
  
```

0036282003405

20					25					30					
Pro	Ile	Pro	Ser	Pro	Phe	Ala	Asp	Gly	Arg	Glu	Gln	Pro	Val	Glu	Val
35				40				45							
Arg	Tyr	Ala	Thr	Ser	Ala	Ala	Ala	Cys	Asp	Met	Leu	Ala	Leu	Ile	Ala
50				55				60							
Asp	Pro	Gln	Val	Gly	Arg	Thr	Leu	Trp	Glu	Ala	Val	Arg	Arg	His	Ala
65				70				75				80			
Arg	Ala	Tyr	Asn	Ala	Thr	Val	Ile	Trp	Tyr	Lys	Ile	Glu	Ser	Gly	Cys
				85				90				95			
Ala	Arg	Pro	Leu	Tyr	Tyr	Met	Glu	Tyr	Thr	Glu	Cys	Glu	Pro	Arg	Lys
100								105				110			
His	Phe	Gly	Tyr	Cys	Arg	Tyr	Arg	Thr	Pro	Pro	Phe	Trp	Asp	Ser	Phe
115								120				125			
Leu	Ala	Gly	Phe	Ala	Tyr	Pro	Thr	Asp	Asp	Glu	Leu	Gly	Leu	Ile	Met
130				135				140							
Ala	Ala	Pro	Ala	Arg	Leu	Val	Glu	Gly	Gln	Tyr	Arg	Arg	Ala	Leu	Tyr
145				150				155				160			
Ile	Asp	Gly	Thr	Val	Ala	Tyr	Thr	Asp	Phe	Met	Val	Ser	Leu	Pro	Ala
				165				170				175			
Gly	Asp	Cys	Trp	Phe	Ser	Lys	Leu	Gly	Ala	Ala	Arg	Gly	Tyr	Thr	Phe
180								185				190			
Gly	Ala	Cys	Phe	Pro	Ala	Arg	Asp	Tyr	Glu	Gln	Lys	Lys	Val	Leu	Arg
195								200				205			
Leu	Thr	Tyr	Leu	Thr	Gln	Tyr	Tyr	Pro	Gln	Glu	Ala	His	Lys	Ala	Ile
210				215				220							
Val	Asp	Tyr	Trp	Phe	Met	Arg	His	Gly	Gly	Val	Val	Pro	Pro	Tyr	Phe
225				230				235				240			
Glu	Glu	Ser	Lys	Gly	Tyr	Glu	Pro	Pro	Pro	Ala	Ala	Asp	Gly	Gly	Ser
				245				250				255			
Pro	Ala	Pro	Pro	Gly	Asp	Asp	Glu	Ala	Arg	Glu	Asp	Glu	Gly	Glu	Thr
260								265				270			
Glu	Asp	Gly	Ala	Ala	Gly	Arg	Glu	Gly	Asn	Gly	Gly	Pro	Pro	Gly	Pro

275 280 285

Glu Gly Asp Gly Glu Ser Gln Thr Pro Glu Ala Asn Gly Gly Ala Glu
 290 300

Gly Glu Pro Lys Pro Gly Pro Ser Pro Asp Ala Asp Arg Pro Glu Gly
 305 310 315 320

Trp Pro Ser Leu Glu Ala Ile Thr His Pro Pro Pro Ala Pro Ala Thr
 325 330 335

Pro Ala Arg Ala
 340

<210> 36
 <211> 1020
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: sequence
 encoding a truncated mature BHV-1 gD

<400> 36

ttgcctacac	ccgcgcgcgc	ggtgacggta	tacgtogacc	cgccggcgta	cccgatgccg	60
cgatacaact	acactgaacg	ctggcacact	accggggcca	taccgtcgcc	cttcgcagac	120
ggccgcgagc	agcccgtcga	ggtgcgctac	gcgacgagcg	cgccggcggtg	cgacatgctg	180
gcgctgatcg	cagaccgcga	ggtggggcgc	acgctgtggg	aagcggtagc	ccggcacgcg	240
cgcgcgtaga	acgccacggt	catatggtac	aagatcgaga	gcgggtgcgc	ccggccgctg	300
tactacatgg	agtacaccga	gtgcgagccc	aggaagcact	ttgggtactg	ccgctaccgc	360
acacccccgt	tttgggacag	cttcctggcg	ggcttcgcct	accccacgga	cgacgagctg	420
ggactgatta	tggcggcgcc	cgcgcggtc	gtcgagggcc	agtaccgacg	cgcgctgtac	480
atcgacggca	cggtcgcccta	tacagatttc	atggtttcgc	tgcgggcggg	ggactgctgg	540
ttctcgaaaac	tcggcgcgccg	tcgcggttac	acctttggcg	cgtgcttccc	ggcccgggat	600
tacgagcaaa	agaaggttct	gcgcctgacg	tatctcacgc	agtactaccc	gcaggaggca	660
cacaaggcca	tagtcgacta	ctggttcattg	cgccacgggg	gcgtcggttc	gccgtatattt	720
gaggagtcga	agggttacga	gccgcgcct	gccgcgatg	gggggtcccc	cgcgccacccc	780
ggcgacgacg	aggcccgcga	ggatgaagg	gagaccgagg	acggggcagc	cgggcgggag	840
ggcaacggcg	gccccccagg	acccgaaggc	gacggcgaga	gtcagacccc	cgaagccaac	900
ggaggcgccg	agggcgagcc	gaaacccggc	cccagccccg	acgcccagccg	ccccgaaggc	960
tggccgagcc	tcgaagccat	cacgcacccc	ccgcccggcc	ccgctacgcc	cgctcgagct	1020

<210> 37
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 6XHIS leader

<400> 37

Met Arg Gly Ser His His His His His Thr Asp Pro His Ala
1 5 10 15

<210> 38

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sequence
encoding 6XHIS leader

<400> 38

atgagaggat ctcacatca ccatcacat acggatccgc atgcc 45

<210> 39

<211> 1017

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: open reading
frame for the 6XHIS leader, truncated mature gD,
and GnRH tetramer encoded by bac-gD:GnRH

<400> 39

atgagcttgc ctacaccgc gccgcgggtg acggtatacg tcgaccgcc gccgtaccgc 60
atgccgggat acaactacac tgaacgctgg cactactacc gccccatacc gtcgcccttc 120
gcagacggcc gcgagcagcc cgtcgaggtg cgctacgcga cgagcgcggc gccgtgcgac 180
atgctggcgc tgatcgaga ccgcaggtg gggcgacgc tgtgggaagc ggtacgccgc 240
cacgcgcgcg cgtacaacgc cacggtcata tggtaacaaga tcgagagcgc gtgcgcccgc 300
ccgctgtact acatggagta caccgagtc gagccagga agcactttgg gtactgccgc 360
taccgcacac ccccgttttg ggacagcttc ctggcgggct tcgcctacc cacggaagac 420
gagctgggac tgattatggc gccgcccgc cggtcgtcg agggccagta ccgacgcgcg 480
ctgtacatcg acggcacggt cgctataca gatttcattg ttctcgtgcc gcccggggac 540
tgctggttct cgaaactcgc cgcggctcgc gggtacacct ttggcgcgtg cttcccgcc 600
cgggattacg agcaaaagaa ggttctgcgc ctgacgtatc tcacgcagta ctaccgcag 660
gaggcacaca aggccatagt cgactactgg ttcatgcgc acgggggcgt cgttcgccgc 720
tattttgagg agtcgaagg ctacgagccg ccgcctgcc ccgatgggg ttccccgcgc 780
ccaccgggcg acgacgagc ccgcgaggt gaaggggaga ccgagacgc gccagccggg 840
cgggagggca acggcggccc ccagagacc gaaggcgacg gcgagagtc gacccccgaa 900

gccaacggag gcgccgaggg cgagccgaaa cccggcccca gccccgacgc cgaccgcccc 960
gaaggctggc cgagcctcga agccatcacg cccccccgc ccgccccgcg tacgcc 1017

<210> 40

<211> 1272

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sequence
encoding a 4GnRH-tmgD-4GnRH fusion protein

<400> 40

gagcactggg catatggtct gcgtccgggt gaacattgga gctacggtct acgccccggg 60
gaacactggg cttatggctt acggccggga gagcattgga gttacggcct ccgtccaggt 120
tccatgagct tgccacacc cgcgcgcggg gtgacgggtat acgtcgaccc gccggcggtac 180
ccgatgccgc gatacaacta cactgaacgc tggcacacta ccggggcccat accgtcgccc 240
ttcgcagacg gccgcgagca gcccgctcag gtgcgctacg cgacgagcgc ggcggcgtgc 300
gacatgctgg cgctgatcgc agaccgcgag gtggggcgca cgctgtggga agcggtacgc 360
cggcacgcgc gcgcgtacaa cgccacggtc atatggtaca agatcgagag cgggtgcgcc 420
cggccgctgt actacatgga gtacaccgag tgcgagccca ggaagcactt tgggtactgc 480
cgctaccgca ccccccggtt ttgggacagc ttcttgccgg gcttcgccta cccacaggac 540
gacgagctgg gactgattat ggccggcgccc gcgcggctcg tcgagggcca gtaccgacgc 600
gcgctgtaca tcgacggcac ggtgcctat acagatttca tggtttcgct gccggccggg 660
gactgctggg tctcgaaact cggcgcggtc cgcggttaca cctttggcgc gtgcttcccg 720
gcccgggatt acgagcaaaa gaaggttctg cgctgacgt atctcacgca gtactacccg 780
caggaggcac acaaggccat agtcgactac tggttcatgc gccacggggg cgtcgttccg 840
cogtattttg aggagtcgaa gggctacgag ccgcccctg ccgccgatgg gggttcccc 900
gcgccacccg gcgacgacga ggcccgcgag gatgaagggg agaccgagga cggggcagcc 960
gggcggggag gcaacggcgg cccccagga cccgaaggcg acggcgagag tcagaccccc 1020
gaagccaacg gaggcgccga gggcgagccg aaaccgggcc ccagccccga cgccgaccgc 1080
cccgaaggct ggccgagcct cgaagccatc acgcaccccc cgcccgcccc cgctacgccc 1140
gctcgagctc cagagcactg gtcatatggg ctgcgtccgg gtgaacattg gagctacggg 1200
ctaogccccg gggaacactg gtcttatggc ttacggccgg gagagcattg gagttacggc 1260
ctccgtccag gt 1272

<210> 41

<211> 1144

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: sequence
encoding a tmgD-4GnRH fusion protein

<400> 41

cttgcctaca cccgcgcgcg ggggtgacggt atacgtcgac ccgcccggcgt acccgatgcc 60
 gcgatacaac tacactgaac gctggcacac taccggggccc ataccgtcgc ctttcgcaga 120
 cgcccgcgag cagcccgctg aggtgcgcta cgcgacgagc gcggcgggcgt gcgacatgct 180
 ggcgctgata gcagacccgc aggtggggcg cacgctgtgg gaagcggtag gccggcacgc 240
 gcgcgcgtac aacgccacgg tcatatggta caagatcgag agcgggtgcg cccggccgct 300
 gtactacatg gagtacaccg agtgcgagcc caggaagcac tttgggtact gccgctaccg 360
 cacacccccg ttttgggaca gcttcctggc gggcttcgcc taccacacgg acgacgagct 420
 gggactgatt atggcggcgc ccgcgcggct cgtcgagggc cagtaccgac gcgcgctgta 480
 catcgacggc acggtcgcct atacagattt catggtttcg ctgcggggcg gggactgctg 540
 gttctcgaaa ctgcgcgcgg ctgcgggta cacctttggc gcgtgcttcc cggcccggga 600
 ttacgagcaa aagaaggttc tgcgcctgac gtatctcacg cagtactacc cgcaggaggc 660
 acacaaggcc atagtcgact actggttcat gcgccacggg ggcgtcgttc cgcgctattt 720
 tgaggagtcg aagggtctacg agccgcgcgc tgccgcgcgat gggggttccc ccgcgccacc 780
 cggcgacgac gaggcccgcg aggatgaagg ggagaccgag gacggggcag ccgggcggga 840
 gggcaacggc ggccccccag gacccgaagg cgacggcgag agtcagaccc ccgaagccaa 900
 cggagggcgc gaggcgagc cgaaacccgg cccagcccc gacgccgacc gccccgaagg 960
 ctggccgagc ctggaagcca tcacgcaccc cccgcccgc cccgctacgc ccgctcgagc 1020
 tccagagcac tggtcatatg gtctgcgtcc gggatgaacat tggagctacg gtctacgccc 1080
 cggggaacac tggctttatg gcttacggcc gggagagcat tggagttacg gcctccgtcc 1140
 aggt 1144

<210> 42

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer P14-S1

<400> 42

ggagctccag agcactggtc ata 23

<210> 43

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer
P14-A138

<400> 43

aaagcttcaa cctggacgga ggcc 24

<210> 44

<211> 215
 <212> PRT
 <213> Actinobacillus pleuropneumoniae

<400> 44

Met	Lys	Lys	Ala	Val	Leu	Ala	Ala	Val	Leu	Gly	Gly	Ala	Leu	Leu	Ala	1	5	10	15
Gly	Ser	Ala	Met	Ala	His	Gln	Ala	Gly	Asp	Val	Ile	Phe	Arg	Ala	Gly	20	25	30	
Ala	Ile	Gly	Val	Ile	Ala	Asn	Ser	Ser	Ser	Asp	Tyr	Gln	Thr	Gly	Ala	35	40	45	
Asp	Val	Asn	Leu	Asp	Val	Asn	Asn	Asn	Ile	Gln	Leu	Gly	Leu	Thr	Gly	50	55	60	
Thr	Tyr	Met	Leu	Ser	Asp	Asn	Leu	Gly	Leu	Glu	Leu	Leu	Ala	Ala	Thr	65	70	75	80
Pro	Phe	Ser	His	Lys	Ile	Thr	Gly	Lys	Leu	Gly	Ala	Thr	Asp	Leu	Gly	85	90	95	
Glu	Val	Ala	Lys	Val	Lys	His	Leu	Pro	Pro	Ser	Leu	Tyr	Leu	Gln	Tyr	100	105	110	
Tyr	Phe	Phe	Asp	Ser	Asn	Ala	Thr	Val	Arg	Pro	Tyr	Val	Gly	Ala	Gly	115	120	125	
Leu	Asn	Tyr	Thr	Arg	Phe	Phe	Ser	Ala	Glu	Ser	Leu	Lys	Pro	Gln	Leu	130	135	140	
Val	Gln	Asn	Leu	Arg	Val	Lys	Lys	His	Ser	Val	Ala	Pro	Ile	Ala	Asn	145	150	155	160
Leu	Gly	Val	Asp	Val	Lys	Leu	Thr	Asp	Asn	Leu	Ser	Phe	Asn	Ala	Ala	165	170	175	
Ala	Trp	Tyr	Thr	Arg	Ile	Lys	Thr	Thr	Ala	Asp	Tyr	Asp	Val	Pro	Gly	180	185	190	
Leu	Gly	His	Val	Ser	Thr	Pro	Ile	Thr	Leu	Asp	Pro	Val	Val	Leu	Phe	195	200	205	
Ser	Gly	Ile	Ser	Tyr	Lys	Phe	210	215											

<210> 45
 <211> 364
 <212> PRT
 <213> Actinobacillus pleuropneumoniae

<400> 45

```

Met Lys Lys Ser Leu Val Ala Leu Thr Val Leu Ser Ala Ala Ala Val
  1             5             10             15

Ala Gln Ala Ala Pro Gln Gln Asn Thr Phe Tyr Ala Gly Ala Lys Ala
      20             25             30

Gly Trp Ala Ser Phe His Asp Gly Ile Glu Gln Leu Asp Ser Ala Lys
      35             40             45

Asn Thr Asp Arg Gly Thr Lys Tyr Gly Ile Asn Arg Asn Ser Val Thr
      50             55             60

Tyr Gly Val Phe Gly Gly Tyr Gln Ile Leu Asn Gln Asp Lys Leu Gly
      65             70             75             80

Leu Ala Ala Glu Leu Gly Tyr Asp Tyr Phe Gly Arg Val Arg Gly Ser
      85             90             95

Glu Lys Pro Asn Gly Lys Ala Asp Lys Lys Thr Phe Arg His Ala Ala
      100            105            110

His Gly Ala Thr Ile Ala Leu Lys Pro Ser Tyr Glu Val Leu Pro Asp
      115            120            125

Leu Asp Val Tyr Gly Lys Val Gly Ile Ala Leu Val Asn Asn Thr Tyr
      130            135            140

Lys Thr Phe Asn Ala Ala Gln Glu Lys Val Lys Thr Arg Arg Phe Gln
      145            150            155            160

Ser Ser Leu Ile Leu Gly Ala Gly Val Glu Tyr Ala Ile Leu Pro Glu
      165            170            175

Leu Ala Ala Arg Val Glu Tyr Gln Trp Leu Asn Asn Ala Gly Lys Ala
      180            185            190

Ser Tyr Ser Thr Leu Asn Arg Met Gly Ala Thr Asp Tyr Arg Ser Asp
      195            200            205

Ile Ser Ser Val Ser Ala Gly Leu Ser Tyr Arg Phe Gly Gln Gly Ala
      210            215            220

```

003700-02090860

Val Pro Val Ala Ala Pro Ala Val Glu Thr Lys Asn Phe Ala Phe Ser
225 230 235 240

Ser Asp Val Leu Phe Ala Phe Gly Lys Ser Asn Leu Lys Pro Ala Ala
245 250 255

Ala Thr Ala Leu Asp Ala Met Gln Thr Glu Ile Asn Asn Ala Gly Leu
260 265 270

Ser Asn Ala Ala Ile Gln Val Asn Gly Tyr Thr Asp Arg Ile Gly Lys
275 280 285

Glu Ala Ser Asn Leu Lys Leu Ser Gln Arg Arg Ala Glu Thr Val Ala
290 295 300

Asn Tyr Ile Val Ser Lys Gly Ala Pro Ala Ala Asn Val Thr Ala Val
305 310 315 320

Gly Tyr Gly Glu Ala Asn Pro Val Thr Gly Ala Thr Cys Asp Lys Val
325 330 335

Lys Gly Arg Lys Ala Leu Ile Ala Cys Leu Ala Pro Asp Arg Arg Val
340 345 350

Glu Val Gln Val Gln Gly Thr Lys Glu Val Thr Met
355 360

<210> 46
<211> 369
<212> PRT
<213> Actinobacillus pleuropneumoniae

<400> 46
Met Lys Lys Ser Leu Val Ala Leu Ala Val Leu Ser Ala Ala Ala Val
1 5 10 15

Ala Gln Ala Ala Pro Gln Gln Asn Thr Phe Tyr Ala Gly Ala Lys Val
20 25 30

Gly Gln Ser Ser Phe His His Gly Val Asn Gln Leu Lys Ser Gly His
35 40 45

Asp Asp Arg Tyr Asn Asp Lys Thr Arg Lys Tyr Gly Ile Asn Arg Asn
50 55 60

Ser Val Thr Tyr Gly Val Phe Gly Gly Tyr Gln Ile Leu Asn Gln Asn
65 70 75 80

Thr Cys Asp Lys Val Lys Gly Arg Lys Ala Leu Ile Ala Cys Leu Ala
340 345 350

Pro Asp Arg Arg Val Glu Val Gln Val Gln Gly Ala Lys Asn Val Ala
355 360 365

Met

003727 3203360